

**Instructions:** This was a 2 hour, closed notes, closed book, and pledged exam.

1. Find and classify all the critical points of

$$f(x, y) = \frac{1}{2}x^2 - xy + \frac{1}{3}y^3.$$

*Solution:* Since  $f$  is differentiable everywhere, the only critical points are the points where the gradient vanishes.  $\nabla f(x, y) = (x - y, y^2 - x) = (0, 0)$  precisely at the points  $(0, 0)$  and  $(1, 1)$ . Note  $\frac{\partial^2 f}{\partial x^2} = 1$ ,  $\frac{\partial^2 f}{\partial y \partial x} = -1$ , and  $\frac{\partial^2 f}{\partial y^2} = 2y$ . Apply the second derivative test:

point	$\frac{\partial^2 f}{\partial x^2}$	$D$	classification
$(0, 0)$	1	-1	saddle
$(1, 1)$	1	1	strict local min

2. Let  $g(x, y) = 2e^{-x} \cos y$ .

- (a) Find the quadratic Taylor polynomial for  $g(x, y)$  around the point  $(0, 0)$ .

$$\begin{aligned} g(0, 0) &= 2 \\ \frac{\partial g}{\partial x} \Big|_{(0,0)} &= -2e^{-x} \cos y \Big|_{(0,0)} = -2 \\ \frac{\partial^2 g}{\partial x^2} \Big|_{(0,0)} &= 2e^{-x} \cos y \Big|_{(0,0)} = 2 \\ \frac{\partial^2 g}{\partial y \partial x} \Big|_{(0,0)} &= -2e^{-x} \sin y \Big|_{(0,0)} = 0 \\ \frac{\partial g}{\partial y} \Big|_{(0,0)} &= -2e^{-x} \sin y \Big|_{(0,0)} = 0 \\ \frac{\partial^2 g}{\partial y^2} \Big|_{(0,0)} &= -2e^{-x} \cos y \Big|_{(0,0)} = -2 \end{aligned}$$

So the quadratic Taylor polynomial for  $g(x, y)$  around  $(0, 0)$  is

$$g(h_1, h_2) \approx 2 - 2h_1 + h_1^2 - h_2^2$$

- (b) Use your answer in part (a) to estimate  $2e^{-0.2} \cos 0.4$ .

We just evaluate  $g(0.2, 0.4) \approx 2 - 2(0.2) + (0.2)^2 - (0.4)^2 = 2 - .4 + .04 - .16 = 1.48$ .  
(The actual value is approximately 1.508202).

3. A tank is in the shape of a half-cylinder of radius 2 and height 3. It is situated in  $\mathbb{R}^3$ , given by the inequalities  $\sqrt{x^2 + y^2} \leq 2$ ,  $y \geq 0$ , and  $0 \leq z \leq 3$ . The temperature at the point  $(x, y, z)$  is given by

$$T(x, y, z) = 2yz^2 \sqrt{x^2 + y^2} \text{ } ^\circ\text{C}.$$

Find the average temperature in the tank.

*Solution:* We describe the tank in cylindrical coordinates as  $0 \leq r \leq 2, 0 \leq \theta \leq \pi$ , and  $0 \leq z \leq 3$ . Recall the formula  $[T]_{av} = \frac{\iiint_W T(x,y,z) dV}{\iiint_W dV}$ .

We use cylindrical coordinates to compute

$$\begin{aligned} \iiint_W T(x,y,z) dV &= \int_0^3 \int_0^\pi \int_0^2 2(r \sin \theta)(z^2)(r) r dr d\theta dz \\ &= 2 \left( \int_0^3 z^2 dz \right) \left( \int_0^\pi \sin \theta d\theta \right) \left( \int_0^2 r^3 dr \right) \\ &= 2(2)(4)(9) = 144 \end{aligned}$$

The denominator (volume of region) is given by the formula  $\text{Vol}(W) = \frac{\pi \cdot 2^2 \cdot 3}{2} = 6\pi$ , or by computing

$$\iiint_W dV = \int_0^3 \int_0^\pi \int_0^2 r dr d\theta dz = \left( \int_0^3 dz \right) \left( \int_0^\pi d\theta \right) \left( \int_0^2 r dr \right) = (3)(\pi)(2) = 6\pi$$

Thus,  $[T]_{av} = \frac{144}{6\pi} = \frac{24}{\pi} \circ C$ .

4. Let  $T$  be the triangle with vertices  $(0, 0)$ ,  $(1, 1)$  and  $(0, 1)$  and let  $f(x, y) = x \sin(y^3)$ .

(a) Find the correct limits of integration to **set up**  $\iint_T f(x, y) dA$  as a double integral

$$\iint f(x, y) dx dy.$$

*Solution:*  $\int_0^1 \int_0^y f(x, y) dx dy$ .

(b) Find the correct limits of integration to **set up**  $\iint_T f(x, y) dA$  as a double integral

$$\iint f(x, y) dy dx.$$

*Solution:*  $\int_0^1 \int_x^1 f(x, y) dy dx$ .

(c) Compute  $\iint_T f(x, y) dA$ .

*Solution:* Use the set up from (a):

$$\begin{aligned} \iint_T f(x, y) dA &= \int_0^1 \int_0^y x \sin(y^3) dx dy \\ &= \int_0^1 \frac{\sin(y^3)}{2} x^2 \Big|_{x=0}^y dy \\ &= \int_0^1 \frac{1}{2} y^2 \sin(y^3) dy \\ &= -\frac{\cos(y^3)}{6} \Big|_{y=0}^1 \\ &= \frac{1 - \cos 1}{6} \end{aligned}$$

5. Find the maximum and minimum values obtained by  $f(x, y) = x + y^2$  on the ellipse  $x^2 + 3y^2 \leq 9$ .

*Solution:* First, find critical points in the interior  $x^2 + 3y^2 < 9$ . Note  $\nabla f(x, y) = (1, 2y)$  is never  $(0, 0)$ , so there are no critical points in the interior.

Second, find critical points on the boundary  $x^2 + 3y^2 = 9$  using Lagrange Multipliers. Our constraint function is  $g(x, y) = x^2 + 3y^2$ . Solve  $\nabla f(x, y) = \lambda \nabla g(x, y)$ , i.e.  $(1, 2y) = \lambda(2x, 6y)$ . The second coordinate gives two possibilities:  $y = 0$  or  $\lambda = 1/3$ . If  $y = 0$ , then  $x = \pm 3$  (from the constraint  $x^2 + 3y^2 = 9$ ). If  $\lambda = 1/3$ , then  $x = 3/2$  (from  $1 = 2\lambda x$ ), and the constraint gives  $y = \pm 3/2$ . There are four critical points to investigate:  $(\pm 3, 0)$  and  $(3/2, \pm 3/2)$ .

$(x, y)$	$f(x, y)$
$(3, 0)$	3
$(-3, 0)$	-3
$(3/2, 3/2)$	15/4
$(3/2, -3/2)$	15/4

Thus the absolute maximum value of  $f$  on boundary is  $15/4$ , and the absolute minimum value on the boundary is  $-3$ .

Since there are no critical points from the interior, these maximum and minimum boundary values are also the maximum and minimum values throughout the entire region.

6. The region  $S$  is cut from a solid ball of radius 1 centered at the origin.  $S$  is the region cut by the inequalities  $z \geq 0$  and  $y \geq x$ . ( $S$  is one-quarter of the entire ball, and contains the point  $(0, 1, 0)$ .)

The mass density of  $S$  at a point  $(x, y, z)$  is given by the function  $\delta(x, y, z) = 30z^2 \text{ kg/m}^3$ .

- (a) Find the total mass of  $S$ .

*Solution:* The total mass is given by  $\iiint_S \delta(x, y, z) dV$ . Note that  $S$  is described in spherical coordinates by  $0 \leq \rho \leq 1$ ,  $\pi/4 \leq \theta \leq 5\pi/4$ , and  $0 \leq \phi \leq \pi/2$ . Thus

$$\begin{aligned}
 \iiint_S \delta(x, y, z) dV &= \int_0^{\pi/2} \int_{\pi/4}^{5\pi/4} \int_0^1 30(\rho \cos \phi)^2 \rho^2 \sin \phi d\rho d\theta d\phi \\
 &= 30 \left( \int_0^{\pi/2} \cos^2 \phi \sin \phi d\phi \right) \left( \int_{\pi/4}^{5\pi/4} d\theta \right) \left( \int_0^1 \rho^4 d\rho \right) \\
 &= 30 \left( \left. \frac{-\cos^3 \phi}{3} \right|_0^{\pi/2} \right) (\pi) \left( \frac{1}{5} \right) \\
 &= 2\pi \text{ kg}
 \end{aligned}$$

- (b) Find the average mass density of  $S$ .

*Solution:* Average mass density is

$$[\delta]_{\text{av}} = \frac{\iiint_S \delta(x, y, z) dV}{\iiint_S dV} = \frac{2\pi \text{ kg}}{\text{Vol}(S)} = \frac{2\pi \text{ kg}}{\pi/3 \text{ m}^3} = 6 \text{ kg/m}^3$$