

## BV Compactness for Maps to a Metric Space

Suppose  $-\infty \leq a < b \leq \infty$ ,  $E$  is a metric space,  $e_0$  is any fixed point of  $E$ , and  $f : (a, b) \rightarrow E$  is Lebesgue measurable.

**Def.** We say  $f$  belongs to  $L^1((a, b)^n, E)$  if  $\int_{(a, b)^n} \text{dist}_E(f(x), e_0) dx < \infty$ .

**Lemma 1.** If  $f_j, f : (a, b)^n \rightarrow E$  are  $\mathcal{L}^n$  measurable and

$$\Lambda_j \equiv \int_{(a, b)^n} \text{dist}_E(f_j(x), f(x)) dx \rightarrow 0 \text{ as } j \rightarrow \infty ,$$

then a subsequence  $f_{j'}$  converges pointwise a.e. to  $f$ .

*Proof :* Choose a subsequence  $f_{j'}$  so that  $\sum_{j=1}^{\infty} \Lambda_j < \infty$ . Then, since

$$\int_{(a, b)^n} \sum_{j=1}^{\infty} \text{dist}_E(f_j(x), f(x)) dx < \infty ,$$

$\sum_{j=1}^{\infty} \text{dist}_E(f_j(x), f(x)) < \infty$  for a.e.  $x \in (a, b)^n$  and  $f_{j'}(x) \rightarrow f(x)$  for all such  $x$ . ■

For a measurable map  $f : (a, b) \rightarrow E$ , we define the *essential variation*

$$\text{ess } V_a^b(f) = \sup \left\{ \sum_{i=1}^m \text{dist}_E(f(t_i), f(t_{i-1})) : a < t_0 < t_1 < \dots < t_m < b, \right.$$

$t_i$  are Lebesgue pts of  $f$  } .

Suppose  $f \in L^1((a, b), \mathbf{R})$  and  $\text{ess } V_a^b(f) < \infty$ . Then  $f$  equals a.e. the difference of the two monotone functions  $\text{ess } V_a^x(f) - [\text{ess } V_a^b(f) - f(x)]$ . It follows that the limit  $\tilde{f}(x) = \lim_{r \downarrow 0} \frac{1}{|\mathbf{B}_r(x)|} \int_{\mathbf{B}_r(x)} f(y) dy$  exists at *all* points in  $(a, b)$  and is continuous except for an atmost countable set where the left and right limits still exist but are different. By Lebesgue's differentiation theorem,  $f = \tilde{f}$  a.e., and so  $\text{ess } V_a^b(f)$  coincides with the classical variation of  $\tilde{f}$  :

$$V_a^b(\tilde{f}) = \sup \left\{ \sum_{i=1}^m \text{dist}_E(\tilde{f}(t_i), \tilde{f}(t_{i-1})) : a < t_0 < t_1 < \dots < t_m < b \right\} .$$

For an open set  $U \subset \mathbf{R}^n$ , recall also the distribution definition:

**Def.**  $f \in BV(U) \Leftrightarrow f \in L^1(U)$  and

$$\|Df\|(U) \equiv \sup \left\{ \int f \text{div } w : w \in \mathcal{C}_0^\infty(U, \mathbf{R}^n), |w| \leq 1 \right\} < \infty .$$

For  $n = 1$  we have:

**Theorem 1.** If  $f \in L^1((a, b))$ , then  $\|Df\|((a, b)) = \text{ess } V_a^b(f)$ .

The proof given in class, used smoothing for both implications and followed Evans-Gariepy, §5.10.1. ■

**Lemma 2.** Suppose  $0 < M < \infty$  and  $f_j : [a, b] \rightarrow [-M, M]$  are monotone increasing. Then a subsequence  $f_{j'}$  converges pointwise off a countable set. Moreover,  $\|f_{j'} - f\|_{L^p} \rightarrow 0$  for any  $p \in [1, \infty)$ .

*Proof :* Suppose  $\mathbf{Q} \cap [a, b] = \{a_1, a_2, \dots\}$ . A subsequence  $f_{\alpha_1(1)}(a_1), f_{\alpha_1(2)}(a_1), \dots$  of the bounded sequence of numbers  $f_1(a_1), f_2(a_1), \dots$  converges to a number  $f(a_1)$ . Inductively, choose a subsequence  $f_{\alpha_j(1)}(a_j), f_{\alpha_j(2)}(a_j), \dots$  of the sequence  $f_{\alpha_{j-1}(1)}(a_j), f_{\alpha_{j-1}(2)}(a_j), \dots$  convergent to a number  $f(a_j)$ .

Let  $j' = \alpha_j(j)$  and  $f(x) = \sup_{a_i < x} f(a_i) = \lim_{\epsilon \downarrow 0} \sup_{x-\epsilon < a_i < x} f(a_i)$ . Then  $f$  is monotone increasing and the set  $Z$  of discontinuities of  $f$  is at most countable. To see that  $\lim_{j \rightarrow \infty} f_{j'}(x) = f(x)$  for any  $x \in (a, b) \setminus Z$ , we choose, for  $\epsilon > 0$ , numbers  $a_i < x < a_{\bar{i}}$  so that  $f(a_{\bar{i}}) - \epsilon < f(x) < f(a_i) + \epsilon$ , and then  $J$  so that

$$|f_{j'}(a_i) - f(a_i)| < \epsilon \text{ and } |f_{j'}(a_{\bar{i}}) - f(a_{\bar{i}})| < \epsilon$$

for  $j \geq J$ . For such  $j$  it follows that

$$f(x) - 2\epsilon < f(a_i) - \epsilon < f_{j'}(a_i) < f_{j'}(x) < f_{j'}(a_{\bar{i}}) < f(a_{\bar{i}}) + \epsilon < f(x) + 2\epsilon.$$

Thus  $|f_{j'}(x) - f(x)| < 2\epsilon$ .

To verify the second conclusion note that  $|f_{j'} - f|^p \leq 2^p M^p$  and apply the Lebesgue dominated convergence theorem. ■

**Corollary 1.** Any sequence of functions  $f_j \in L^1((a, b))$  with

$$M \equiv \sup_j \int_a^b |f_j| dx + \|Df_j\|((a, b)) < \infty$$

contains a subsequence  $f_{j'}$  convergent pointwise a.e. and in  $L^1$  to a function  $f$  with  $\int_a^b |f| dx + \|Df\|((a, b)) \leq M$ .

*Proof :* For the normalized functions  $\tilde{f}_j$  we have the sup bound

$$\begin{aligned} |\tilde{f}_j| &\leq |b - a| V_a^b \tilde{f}_j + \inf |\tilde{f}_j| \\ &\leq |b - a| \|Df_j\|((a, b)) + \frac{1}{|b - a|} \int_a^b |f_j| \leq M(|b - a| + |b - a|^{-1}). \end{aligned}$$

We may as above then write  $\tilde{f}_j$  as the difference  $g_j - h_j$  of two uniformly bounded monotone increasing functions. Applying Lemma 2 to  $g_j$  and  $h_j$  gives the Corollary. ■

Recall that a metric space  $E$  is *weakly separable* if there is a sequence of functions  $\phi_i : E \rightarrow \mathbf{R}$  with  $\text{Lip}(\phi_i) \leq 1$  so that

$$\text{dist}_E(x, y) = \inf_i |\phi_i(x) - \phi_i(y)| \text{ for all } x, y \in E .$$

A separable metric space  $E$  is weakly separable as one sees by taking  $\phi_i(x) = \text{dist}_E(x, e_i)$  for some countable dense subset  $\{e_i\}$  of  $E$ . Moreover,  $E$  is weakly separable if and only if there is a distance preserving embedding of  $E$  into

$$\ell^\infty = \{(a_1, a_2, \dots) : \sup_i |a_i| < \infty\} .$$

With  $\phi_i$  as above, one such embedding is

$$\iota(x) = (\phi_1(x) - \phi_1(e_0), \phi_2(x) - \phi_2(e_0), \dots)$$

where  $e_0$  is any given point of  $E$ .

We also say  $E$  is *boundedly compact* if every closed ball  $\overline{\mathbf{B}}_R(e_0) = \{x : \text{dist}_E(x, e_0) \leq R\}$  is compact for  $0 < R < \infty$ . This implies that  $E$  is locally compact and complete.

**Corollary 2.** (BV compactness for  $n = 1$ ) *Suppose  $E$  is a boundedly compact weakly separable metric space,  $e_0 \in E$ , and  $0 < a < b < \infty$ . Any sequence of functions  $f_j \in L^1((a, b), E)$  with*

$$M \equiv \sup_j \int_a^b \text{dist}_E(f_j(x), e_0) dx + \text{ess } V_a^b(f_j) < \infty$$

*contains a subsequence  $f_{j'}$  convergent pointwise a.e. and in  $L^1$  to a function  $f$  with  $\int_a^b \text{dist}_E(f(x), e_0) dx + \text{ess } V_a^b(f) \leq M$ .*

*Proof:* We may assume  $E$  is a boundedly compact subset of  $\ell^\infty$  and write  $f_j = (f_j^1, f_j^2, \dots)$ . We apply Corollary 1 first to the sequence  $(f_1^1, f_2^1, \dots)$  to obtain a subsequence  $f_{\alpha_1(j)}^1$  that is  $L^1$  and pointwise a.e. convergent to some  $f^1$ , then inductively to the sequence  $f_{\alpha_{k-1}(j)}^k$  to obtain a subsequence  $f_{\alpha_k(j)}^k$  convergent to some  $f^k$ . We conclude that, for the diagonal sequence  $f_{j'} = f_{\alpha_j(j)}$ , each  $f_{j'}^k$  is  $L^1$  and pointwise a.e. convergent to  $f^k$  as  $j \rightarrow \infty$ . But, for the convergence of the functions  $\|f_{j'}(x) - f(x)\|_{\ell^\infty}$ , we still need to show that the rates of the convergences of  $f_{j'}^k(x)$  to  $f^k(x)$  are uniform independent of  $k$  and that the limit  $f(x) \in E$  for almost every  $x \in (a, b)$ . For this purpose we will show that, for a.e.  $x \in (a, b)$ , any subsequence  $j''$  of  $j'$  contains a subsequence  $j'''$  so that  $f_{j'''}(x) \rightarrow f(x)$ . First Fatou's Lemma gives that, for a.e.  $x \in (a, b)$ ,  $\liminf_{j'' \rightarrow \infty} \text{dist}_E(f_{j''}(x), e_0) < \infty$ . So there is, by the bounded compactness assumption, a subsequence  $j'''$  of  $j''$  (depending on  $x$ ) and a limit point  $e \in E$  so that  $\|f_{j'''}(x) - e\|_{\ell^\infty} \rightarrow 0$  as  $j \rightarrow \infty$ . But the previous convergences of the components  $f_{j'}^k$  show that  $e = (e^1, e^2, \dots) = (f^1(x), f^2(x), \dots) = f(x)$ .

Thus we obtain the desired pointwise a.e. convergence. The  $L^1$  convergence then follows by Lebesgue dominated convergence as in the proof of Corollary 1, and one readily checks that

$$\int_a^b \|f(x)\|_{\ell^\infty} dx + \text{ess } V_a^b(f) \leq M .$$

■

Now we turn again to functions of  $n$  variables. For  $x \in \mathbf{R}^n$ ,  $k \in \{1, \dots, n\}$ , and  $f : (a, b)^n \rightarrow E$ , we define

$$\hat{x}_k = (x_1, \dots, x_{k-1}, x_{k+1}, \dots, x_n), \quad f_{(k)}(\hat{x}_k, x_k) = f(x) .$$

Then we have:

**Lemma 3.** For  $f \in L^1((a, b)^n, E)$  with  $E$  weakly separable, the mapping

$$y \mapsto \text{ess } V_a^b f_{(k)}(y, \cdot)$$

is  $\mathcal{L}^{n-1}$  measurable.

**Theorem 2.** Suppose  $f \in L^1((a, b)^n)$ . Then

$$\|Df\|((a, b)^n) \leq \sum_{k=1}^n \int_{(a, b)^n} \text{ess } V_a^b f_{(k)}(y, \cdot) dy \leq n \|Df\|((a, b)^n) .$$

The proofs given in class followed Evans-Gariepy, §5.10.2. ■

Based on the above it is reasonable to say that a function  $f \in L^1((a, b)^n, E)$  belongs to  $BV((a, b)^n, E)$  if the *variation on lines*

$$VL(f) \equiv \sum_{k=1}^n \int_{(a, b)^n} \text{ess } V_a^b f_{(k)}(y, \cdot) dy < \infty$$

and prove the following:

**Theorem 3.** (BV compactness) Suppose  $E$  is a boundedly compact weakly separable metric space,  $e_0 \in E$ , and  $0 < a < b < \infty$ . Any sequence of functions  $f_j \in L^1((a, b)^n, E)$  with

$$M \equiv \sup_j \int_{(a, b)^n} \text{dist}_E(f_j(x), e_0) dx + VL(f_j) < \infty$$

contains a subsequence  $f_{j'}$  convergent pointwise a.e. and in  $L^1$  to a function  $f$  with

$$\int_{(a, b)^n} \text{dist}_E(f(x), e_0) dx + VL(f) \leq M$$

.

*Proof* : We argue by induction on  $n$ . The case  $n = 1$  follows from Corollary 2. Assuming the theorem true for dimensions less than  $n$ , we will use the metric space

$$\tilde{E} \equiv \{f \in L^1((a, b)^{n-1}, E) : \int_{(a, b)^{n-1}} \text{dist}_E(f(x), e_0) dx + VL(f) \leq M\} .$$

One easily checks that  $\tilde{E}$  is weakly separable by again viewing  $E$  as a boundedly compact subset of  $\ell^\infty$ . The inductive assumption guarantees that  $\tilde{E}$  is compact, hence boundedly compact.

For each  $k \in \{1, \dots, n\}$ , Fubini's theorem implies that, for a.e.  $t \in \mathbf{R}$ , each function  $f_{j(k)}(\cdot, t) \in \tilde{E}$  and that the map  $t \mapsto f_{j(k)}(\cdot, t)$  belongs to  $L^1((a, b), \tilde{E})$  with the  $L^1$  norm uniformly bounded by  $M$ . Also for  $a \leq s < t \leq b$ ,

$$\text{dist}_{\tilde{E}}(f_{j(k)}(\cdot, s), f_{j(k)}(\cdot, t)) = \int_{(a, b)^{n-1}} \text{dist}_E(f_{j(k)}(y, s), f_{j(k)}(y, t)) dy$$

so that

$$\text{ess } V_a^b f_{j(k)}(\cdot, \cdot) = \int_{(a, b)^{n-1}} \text{ess } V_a^b f_{j(k)}(y, \cdot) dy \leq M .$$

The case  $n = 1$  now gives  $L^1$  convergence of a subsequence  $f_{j'(k)}(\cdot, t)$  to a function  $g_k \in BV((a, b), \tilde{E})$ . We obtain  $g_1, g_2, \dots, g_n$  by taking consecutive subsequences. The compatibility condition of the approximating functions

$$f_{j(k)}(x'_k, x_k) = f_j(x_1, \dots, x_n) = f_{j(l)}(x'_l, x_l)$$

for  $k, l \in \{1, \dots, n\}$  along with Lemma 1 implies the compatibility of these limit functions

$$g_k(x'_k, x_k) = g_l(x'_l, x_l)$$

which implies, using Fubini's Theorem, the existence of a function well-defined by

$$f(x_1, \dots, x_n) = g_k(x'_k, x_k)$$

for all  $k = 1, \dots, n$  and almost all  $x \in (a, b)^n$ . Also one has, by Fubini's theorem the  $L^1$  convergence

$$\int_{(a, b)^n} \text{dist}_E(f_{j'}, f) dx = \int_a^b \text{dist}_{\tilde{E}}(f_{j'(k)}(\cdot, t), f_{(k)}(\cdot, t)) dt \rightarrow 0$$

as  $j \rightarrow \infty$ . By Lemma 1, a subsequence of the  $f_{j'}$  also converges pointwise a.e. to  $f$ . Measurability is a consequence of the pointwise convergence. One readily verifies the estimate

$$\int_{(a, b)^n} \text{dist}_E(f(x), e_0) dx + VL(f) \leq M$$

.

■