

1 Homework #10

1. Consider the functional with Lagrangian

$$F(x, u, p) = \sqrt{x^2 + u^2} \sqrt{1 + p^2}.$$

- (a) Find the Hamiltonian and the associated Hamiltonian system.

Solution Find that $\pi = \frac{\partial F}{\partial p} = \frac{p\sqrt{x^2+u^2}}{\sqrt{1+p^2}}$, so that (after simplifying), the Hamiltonian is

$$H(x, u, \pi) = \pi p - F(x, u, \pi) = -\sqrt{x^2 + u^2 - \pi^2}.$$

The Hamiltonian system is

$$\begin{aligned} \frac{du}{dx} &= \frac{\partial H}{\partial \pi} = \frac{\pi}{\sqrt{x^2 + u^2 - \pi^2}}, \\ \frac{d\pi}{dx} &= -\frac{\partial H}{\partial u} = \frac{u}{\sqrt{x^2 + u^2 - \pi^2}}. \end{aligned}$$

- (b) Show that $u^2 - \pi^2$ is a conserved quantity.

Solution Using the chain rule,

$$\frac{d}{dx}(u^2 - \pi^2) = 2u \frac{du}{dx} - 2\pi \frac{d\pi}{dx}.$$

From part (a), this derivative is zero, so $u^2 - \pi^2$ is indeed conserved.

- (c) Find a solution to the Hamilton-Jacobi equation with the form

$$S(x, u) = \frac{1}{2} (Ax^2 + 2Bxu + Cu^2).$$

Solution Using $S_x = Ax + Bu$ and $S_u = Bx + Cu$, the Hamilton-Jacobi equation becomes

$$0 = S_x + H(x, u, S_u) = Ax + Bu - \sqrt{x^2 + u^2 - (Bx + Cu)^2}.$$

Solve this to find that $A = -C$ and $B = \sqrt{1 - A^2}$. A general solution of the given form is then

$$S(x, u) = \frac{1}{2} (Ax^2 + \sqrt{1 - A^2}xu - Au^2).$$

- (d) Show that the solutions are given implicitly in the form

$$(x^2 - u^2) \cos \alpha + 2ux \sin \alpha = C,$$

where α and C are constants.

Solution The extremals will satisfy $S_A = \frac{1}{2} (x^2 - u^2 - \frac{2A}{\sqrt{1-A^2}}ux)$ constant. With $A = \sin \alpha$, find that $x^2 - u^2 - \frac{\sin \alpha}{\cos \alpha}ux$ is constant, so multiplying this by the constant $\cos \alpha$ finishes the problem.

2. Consider a particle with mass m in \mathbb{R}^3 moving under the influence of the potential V . Of course, the Lagrangian is

$$L(x, y, z, p, q, r) = \frac{m}{2}(p^2 + q^2 + r^2) - V(x, y, z).$$

- (a) Write the Lagrangian in polar coordinates r , θ , and ϕ , defined by $x = r \sin \phi \cos \theta$, $y = r \sin \phi \sin \theta$, and $z = r \cos \phi$.

Solution By calculations similar to Homework 3 Exercise 3, find that

$$p^2 + q^2 + r^2 = \dot{x}^2 + \dot{y}^2 + \dot{z}^2 = \dot{r}^2 + r^2 \dot{\phi}^2 + r^2 \sin^2 \phi \dot{\theta}^2.$$

The new Lagrangian is then

$$L(r, \theta, \phi, \dot{r}, \dot{\theta}, \dot{\phi}) = \frac{m}{2}(\dot{r}^2 + r^2 \dot{\phi}^2 + r^2 \sin^2 \phi \dot{\theta}^2) - V(r \sin \phi \cos \theta, r \sin \phi \sin \theta, r \cos \phi).$$

- (b) What is the corresponding Hamiltonian?

Solution Find that

$$\pi = \frac{\partial L}{\partial \dot{r}} = m\dot{r}, \quad \sigma = \frac{\partial L}{\partial \dot{\phi}} = mr^2 \dot{\phi}, \quad \text{and} \quad \rho = \frac{\partial L}{\partial \dot{\theta}} = mr^2 \sin^2 \phi \dot{\theta}.$$

By definition, the Hamiltonian is

$$H(r, \theta, \phi, \pi, \rho, \sigma) = \pi \dot{r} + \sigma \dot{\phi} + \rho \dot{\theta} - L(r, \theta, \phi, \pi, \rho, \sigma),$$

which with the above, simplifies to

$$H(r, \theta, \phi, \pi, \rho, \sigma) = \frac{1}{2m} \left(\pi^2 + \frac{\sigma^2}{r^2} + \frac{\rho^2}{r^2 \sin^2 \phi} \right) + V(r \sin \phi \cos \theta, r \sin \phi \sin \theta, r \cos \phi).$$

- (c) Assuming that $V = -\frac{k}{r}$, reduce the solution of the Hamilton-Jacobi equation to the evaluation of integrals.

Solution The new Hamilton-Jacobi equation is

$$S_t + H(r, \theta, \phi, S_r, S_\theta, S_\phi) = 0$$

$$S_t + \frac{1}{2m} \left(S_r^2 + \frac{S_\theta^2}{r^2 \sin^2 \phi} + \frac{S_\phi^2}{r^2} \right) - \frac{k}{r} = 0.$$

Assume a separable solution

$$S(t, r, \theta, \phi) = f(t) + g(r, \theta, \phi).$$

There must exist a constant α such that

$$f'(t) = -\alpha, \quad \text{and}$$

$$\frac{1}{2m} \left(\left(\frac{\partial g}{\partial r} \right)^2 + \frac{1}{r^2 \sin^2 \phi} \left(\frac{\partial g}{\partial \theta} \right)^2 + \frac{1}{r^2} \left(\frac{\partial g}{\partial \phi} \right)^2 \right) - \frac{k}{r} = \alpha. \quad (1)$$

It is clear that $f(t) = -\alpha t$.

Assume a separable solution

$$g(r, \theta, \phi) = h(r) + w(\theta, \phi).$$

Multiply Equation (1) by r^2 to find that

$$-\alpha r^2 + \frac{1}{2m} \left(r^2 h'(r)^2 + \frac{1}{\sin^2 \phi} w_\theta^2 + w_\phi^2 \right) - kr = 0.$$

With some more simplifying, there must exist a constant β such that

$$\frac{1}{2m} (h'(r)^2 - \alpha) r^2 - kr = -\beta, \text{ and} \quad (2)$$

$$\frac{1}{2m} \left(\frac{1}{\sin^2 \phi} w_\theta^2 + w_\phi^2 \right) = \beta. \quad (3)$$

Assume a separable solution

$$w(\theta, \phi) = u(\theta) + v(\phi).$$

Then from Equation (3),

$$\frac{1}{2m} (u'(\theta)^2 + \sin^2 \phi v'(\phi)^2) - \beta \sin^2 \phi = 0.$$

There exists a constant γ such that

$$\frac{1}{2m} u'(\theta)^2 = -\gamma \text{ and} \quad (4)$$

$$\sin^2 \phi \left(\frac{1}{2m} v'(\phi)^2 - \beta \right) = \gamma. \quad (5)$$

Now to gather up the integrals.

From Equation (2),

$$h(r) = \int \sqrt{\frac{2m}{r^2} (kr - \beta) + \alpha} dr.$$

From Equation (4),

$$u(\theta) = \sqrt{-2m\gamma}\theta.$$

From Equation (5),

$$v(\phi) = \int \sqrt{2m \left(\frac{\gamma}{\sin^2 \phi} + \beta \right)} d\phi.$$

With these three integrals, the solution to the Hamilton-Jacobi equation is $S(t, r, \theta, \phi) = -\alpha t + h(r) + u(\theta) + v(\phi)$.

3. Calculate the second variation of the functional

$$\mathcal{F} = \int_a^b F(x, u(x), u'(x)) dx$$

under the following assumptions.

- (a) $F(x, u, p)$ does not depend on u .

Solution By definition,

$$\delta^2 \mathcal{F}(u, \phi) = \int_a^b (F_{uu}\phi^2 + 2F_{up}\phi\phi' + F_{pp}\phi'^2) dx.$$

With no u dependence,

$$\delta^2 \mathcal{F}(u, \phi) = \int_a^b F_{pp}\phi'^2 dx.$$

- (b) $F(x, u, p)$ does not depend on p .

Solution With no p dependence,

$$\delta^2 \mathcal{F}(u, \phi) = \int_a^b F_{uu}\phi^2 dx.$$

4. Suppose $F(x, u, p) = A(x)u + B(x)p$, where A and B are functions depending only on x . Compute the first and second variations.

Solution The first variation is

$$\delta \mathcal{F}(u, \phi) = \int_a^b (F_u\phi + F_p\phi') dx.$$

It is clear that

$$\delta \mathcal{F}(u, \phi) = \int_a^b (A(x)\phi + B(x)\phi') dx,$$

and

$$\delta^2 \mathcal{F}(u, \phi) = 0.$$

5. Suppose that $F(x, u, p) = A(x) + B(x)u^2 + 2C(x)up + D(x)p^2$. Compute the first and second variations.

Solution Find that

$$F_u = 2B(x)u + 2C(x)p$$

$$F_p = 2C(x)u + 2D(x)p$$

$$F_{uu} = 2B(x)$$

$$F_{up} = 2C(x)$$

$$F_{pp} = 2D(x).$$

Then

$$\delta\mathcal{F}(u, \phi) = \int_a^b (2B(x)u + 2C(x)p)\phi + 2C(x)u + 2D(x)p)\phi' dx,$$

and

$$\delta^2\mathcal{F}(u, \phi) = 2 \int_a^b (B(x)\phi^2 + 2C(x)\phi\phi' + D(x)\phi'^2) dx.$$