

# Math 211

Lecture #3

Solutions to Differential Equations

August 30, 2002

## Differential Equation:

An equation involving an unknown function and one or more of its derivatives, in addition to the independent variable.

- Example:  $y' = \frac{dy}{dt} = 2ty$
- General equation:  $y' = \frac{dy}{dt} = f(t, y)$
- $t$  is the *independent variable*.
- $y = y(t)$  is the *unknown function*.
- $y' = 2ty$  is of *order 1*.

[Return](#)

## Solutions to Differential Equations

The general first order equation is

$$y' = f(t, y).$$

A *solution* is a function  $y(t)$ , defined for  $t$  in an interval, which is differentiable at each point and satisfies

$$y'(t) = f(t, y(t))$$

for every point  $t$  in the interval.

[Return](#)

### Example: $y' = 2ty$

Is  $y(t) = e^{t^2}$  a solution?

- By substitution  $y'(t) = 2ty(t)$ , so  $y(t) = e^{t^2}$  is a solution.

Is  $y(t) = e^t$  a solution?

- By substitution  $y'(t) \neq 2ty(t)$ , so  $y(t) = e^t$  is *not* a solution to the equation  $y' = 2ty$ .

Verification by substitution is always available.

[Return](#)

[Definition of ODE](#)

### More about Solutions

- A solution is a function. What is a function?
  - ♦ An exact, algebraic formula (e.g.,  $y(t) = e^{t^2}$ ).
  - ♦ A convergent power series.
  - ♦ The limit of a sequence of functions.
- An ODE is a function generator.
- Two of the themes of the course are aimed at those solutions for which there is no exact formula.

[Definition of solution](#)

[Definition of ODE](#)

### An ODE is a Function Generator

Example:  $y' = y^2 - t$ ,  $y(0) = 0$

- There is no solution to this IVP which can be given using a formula.
- Nevertheless, there is a solution. We can find as many terms in the power series for  $y(t)$  as we want.

$$y(t) = -\frac{1}{2}t^2 + \frac{1}{20}t^5 - \frac{1}{160}t^8 + \dots$$

## Particular and General Solutions

For the equation  $y' = 2ty$

- $y(t) = \frac{1}{2}e^{t^2}$  is a solution. It is a *particular solution*.
- $y(t) = Ce^{t^2}$  is a solution for any constant  $C$ . This is a *general solution*.

General solutions contain arbitrary constants. Particular solutions do not.

[Return](#)

## Initial Value Problem (IVP)

A differential equation & an initial condition.

- Example: Find  $y(t)$  with  $y' = -2ty$  with  $y(0) = 4$ .
- General solution:  $y(t) = Ce^{-t^2}$ .
- Plug in the initial condition:

$$y(0) = 4,$$

$$Ce^0 = 4,$$

$$C = 4$$

Solution to the IVP:  $y(t) = 4e^{-t^2}$ .

[Return](#)

## Normal Form of an Equation

The first order differential equation

$$y' = f(t, y)$$

is said to be in *normal form*.

- Example: The differential equation  $(1 + t^2)y' + y^2 = t^3$  is not in normal form.
- Solve for  $y'$  to put the equation into normal form:

$$y' = \frac{t^3 - y^2}{1 + t^2}$$

- Many statements about differential equations require the equation to be in normal form.

## Interval of Existence

The largest interval over which a solution can exist.

- Example:  $y' = -2ty$ ,  $y(0) = 4$ .
  - ♦ The interval of existence is  $\mathbf{R} = (-\infty, \infty)$ .
- Example:  $y' = 1 + y^2$  with  $y(0) = 1$ .
  - ♦ General solution:  $y(t) = \tan(t + C)$
  - ♦ Initial Condition:  $y(0) = 1 \Rightarrow y(t) = \tan(t + \pi/4)$
  - ♦ The solution exists and is continuous for  $-\pi/2 < t + \pi/4 < \pi/2$ .
  - ♦ The interval of existence is  $-3\pi/4 < t < \pi/4$ .

[Initial value problem](#)

[Return](#)

## Geometric Interpretation of


$$y' = f(t, y)$$

If  $y(t)$  is a solution, and  $y(t_0) = y_0$ , then

$$y'(t_0) = f(t_0, y(t_0)) = f(t_0, y_0).$$

- The slope to the graph of  $y(t)$  at the point  $(t_0, y_0)$  is given by  $f(t_0, y_0)$ .
- Imagine a small line segment attached to each point of the  $(t, y)$  plane with the slope  $f(t, y)$ .

## The Direction Field



## Autonomous Equations

- General equation:  $\frac{dy}{dt} = f(t, y)$
- Autonomous equation:  $\frac{dy}{dt} = f(y)$
- Examples:
  - ♦  $\frac{dy}{dt} = t - y^2$  is not autonomous.
  - ♦  $\frac{dy}{dt} = y(1 - y)$  is autonomous.

In an *autonomous equation* the right-hand side has no explicit dependence on the independent variable.

[Return](#)

## Equilibrium Points

- An *equilibrium point* for the *autonomous equation*  $\frac{dy}{dt} = f(y)$  is a point  $y_0$  such that  $f(y_0) = 0$ .
- Corresponding to the equilibrium point  $y_0$  there is the constant *equilibrium solution*  $y(t) = y_0$ .
- Example:  $\frac{dy}{dt} = y(2 - y)/3$  is an autonomous equation.
  - ♦ The equilibrium points are  $y_0 = 0$  or  $2$ .
  - ♦ The corresponding equilibrium solutions are  $y(t) = 0$  and  $y(t) = 2$ .

[Return](#)

## Between Equilibrium Points

- $\frac{dy}{dt} = f(y) > 0 \Rightarrow y(t)$  is increasing.
- $\frac{dy}{dt} = f(y) < 0 \Rightarrow y(t)$  is decreasing.
- The graphs of solutions to first order equations cannot cross (uniqueness theorem).
- Example:  $\frac{dy}{dt} = y(2 - y)/3$

[Equilibrium point](#)