

UNIFORM BOUNDS FOR WEIL-PETERSSON CURVATURES

MICHAEL WOLF & YUNHUI WU

ABSTRACT. We find bounds for Weil-Petersson holomorphic sectional curvature, and the Weil-Petersson curvature operator in several regimes, that do not depend on the topology of the underlying surface. Among other results, we show that the minimal (most negative) eigenvalue of the curvature operator at any point in the Teichmüller space $\text{Teich}(S_g)$ of a closed surface S_g of genus g is uniformly bounded away from zero. Restricting to a thick part of $\text{Teich}(S_g)$, we show that the minimal eigenvalue is uniformly bounded below by an explicit constant which does not depend on the topology of the surface but only on the given bound on injectivity radius. We also show that the minimal Weil-Petersson holomorphic sectional curvature of a sufficiently thick hyperbolic surface is comparable to -1 .

1. INTRODUCTION

Let S_g be a closed surface of genus g where $g > 1$, and $\text{Teich}(S_g)$ be the Teichmüller space of S_g . Endowed with the Weil-Petersson metric, the Teichmüller space $\text{Teich}(S_g)$ is Kähler ([1]), incomplete ([7, 33]), geodesically complete ([35]) and negatively curved ([30, 34]). Tromba [30] and Wolpert [34] found a formula for the Weil-Petersson curvature tensor, which has been applied to study a variety of curvature properties of $\text{Teich}(S_g)$ over the past several decades. (See also [13, 16, 28] for alternative proofs of the curvature formula.) In their papers, they deduced from their formula that the holomorphic sectional curvatures are bounded above by a negative number which only depends on the genus of the surface, confirming a conjecture of Royden.

We focus in this paper on bounds for Weil-Petersson curvatures that are uniform across Teichmüller spaces, in the sense that the bounds do not depend on the genus g of the surface S_g .

Naturally, there are a number of appealing notions of curvature of a Riemannian manifold. In particular, while we focus on the holomorphic sectional curvatures in the second part of the paper, in the first part of this paper we study the Weil-Petersson curvature operator

$$\tilde{Q} : \wedge^2 T_X \text{Teich}(S_g) \rightarrow \wedge^2 T_X \text{Teich}(S_g),$$

1991 *Mathematics Subject Classification.* 30F60, 53C21, 32G15.

Key words and phrases. Teichmüller space, Weil-Petersson metric, Curvature operator, Holomorphic sectional curvature.

an endomorphism of the $(3g-3)(6g-7)$ -dimensional exterior wedge product space $\wedge^2 T_X \text{Teich}(S_g)$ of the tangent space $T_X \text{Teich}(S_g)$ at $X \in \text{Teich}(S_g)$. Of course, one popular (and important) way to interpret the “curvature” of a manifold is through its sectional curvatures: while these arise as diagonal elements of the curvature operator \tilde{Q} , our interest in the first part of the paper is on the full operator \tilde{Q} , at least in terms of estimating its spectrum.

In [40], by applying the Tromba-Wolpert Weil-Petersson curvature formula, one of us characterized the zero level set of the curvature operator \tilde{Q} and showed that the Riemannian curvature operator of the Weil-Petersson metric is non-positive definite; in particular, some of the eigenvalues of \tilde{Q} are zero and some are negative. In the first part of this paper, we develop some results for the negative eigenvalues, especially as the genus of the surfaces represented in the moduli space becomes arbitrarily large. We survey these in the next four subsections.

In the second part of the paper – described in more detail in the fifth subsection of this introduction – we turn our attention to holomorphic sectional curvatures. We find, for hyperbolic surfaces X_g of genus g of sufficiently large injectivity radius, that we may exhibit holomorphic sections whose curvatures are uniformly bounded away from zero; here the bound does not depend on the genus of the surface but only on the lower bound we have chosen for the injectivity radius.

1.1. Uniform lower bounds on the norm of the curvature operator.

There are recent suggestions that as the genus g of S_g grows large, some regions in the Teichmüller space $\text{Teich}(S_g)$ should become increasingly flat. We show that, from the point of view of the full curvature operator, this is not true: on any sequence of surfaces with genus tending to infinity, at least one of the eigenvalues of \tilde{Q} does not tend to zero. In particular, let $\lambda_{\min}(X)$ denote the minimal, i.e. the most negative, eigenvalue of \tilde{Q} at X . Our first result is

Theorem 1.1. *For any $X \in \text{Teich}(S_g)$ and let $Sca(X)$ be the scalar curvature at X , then*

$$\lambda_{\min}(X) \leq \frac{-1}{2\pi} < 0.$$

More precisely,

$$\lambda_{\min}(X) \leq \frac{2Sca(X)}{9(g-1)}.$$

Because this eigenvalue is the maximum of values of \tilde{Q} applied to (unit) elements of $\wedge^2 T_X \text{Teich}(S_g)$, it is enough to estimate \tilde{Q} on a carefully chosen element of $\wedge^2 T_X \text{Teich}(S_g)$ which yields the bound.

1.2. Uniform upper bounds on the norm of the curvature operator on the thick locus.

It is clear that $\lambda_{\min}(X)$ is less than or equal to the

sectional curvatures at X along arbitrary planes. However, it is well-known that the sectional curvatures of $\text{Teich}(S_g)$ along certain planes could be arbitrarily negative. One specific example is the direction along which a nontrivial simple closed curve of S is pinching to zero. Then that sectional curvature, along the holomorphic plane defined by the pinching direction, tends to negative infinity as the curve goes to zero. More precisely, we let α be a nontrivial simple closed curve on S and $l_\alpha(X)$ be the length of the closed geodesic in X representing α . Consider the gradient $\lambda_\alpha := \text{grad}(l_\alpha^{\frac{1}{2}})$ whose magnitude approaches $\frac{1}{\sqrt{4\pi}}$ as l_α goes to zero. Wolpert in [39] proved that the sectional curvatures along holomorphic planes spanned by λ_α behave as $\frac{-3}{\pi l_\alpha} + O(l_\alpha)$, which goes to negative infinity as α pinches to zero. This refined an earlier estimation of Huang [9]. For more details, one can see Corollary 16 in [39]. Thus, in general we do not have any uniform lower bounds for $\lambda_{\min}(X)$ over $\text{Teich}(S_g)$.

On the other hand, the Mumford compactness theorem [20] implies that the thick part of the moduli space is compact. Since the mapping class group acts on $\text{Teich}(S_g)$ by isometries, the Weil-Petersson curvature operator \tilde{Q} , restricted to the thick part of $\text{Teich}(S_g)$, is bounded. In particular the minimal eigenvalue $\lambda_{\min}(X)$ has a lower bound when X runs over the thick part of $\text{Teich}(S_g)$. Our next result implies that the lower bound may be taken to be uniform, independent of the topology of the surface.

Theorem 1.2. *Given an $\epsilon > 0$, let $\text{Teich}(S_g)^{\geq \epsilon}$ be the ϵ -thick part of $\text{Teich}(S_g)$. Then, there exists a constant $B(\epsilon) > 0$, depending only on ϵ , such that*

$$\lambda_{\min}(X) \geq -B(\epsilon), \quad \forall X \in \text{Teich}(S_g)^{\geq \epsilon}.$$

The argument will present an explicit constant $B(\epsilon)$ whose asymptotics as $\epsilon \rightarrow 0, \infty$ we will exploit in later sections of the paper. In addition, a direct corollary is that the sectional curvature of the Weil-Petersson metric, restricted on the thick part of the moduli space, is uniformly bounded from below, a result due originally to Huang [10].

1.3. Uniform pinched bounds on the norm of the curvature operator on sequences of increasingly thick surfaces.

The subject of the asymptotic geometry of \mathbb{M}_g , the moduli space of S_g , as g tends to infinity, has recently become quite active: see for example Mirzakhani [21, 22, 23, 24] and Cavendish-Parlier [6] (results obtained by refining Brock's [5] quasi-isometry of $\text{Teich}(S_g)$ to the pants graph). In terms of curvature bounds, by combining the results in Wolpert [34] and Teo [29], we may see that, restricted on the thick part of the moduli space, the scalar curvature is comparable to $-g$ as g goes to infinity. In the sixth section of this paper, we study the asymptotic behaviors of certain eigenvalues of \tilde{Q} as the genus goes to infinity.

We recall for context that Buser and Sarnak [4] proved that there exists a family of closed hyperbolic surfaces X_k of genus g_k with $g_k \rightarrow \infty$ as $k \rightarrow \infty$ and $\text{inj}(X_k) \geq \frac{2 \ln g_k}{3}$. Now, we noted above that the constant $B(\epsilon)$ in Theorem 1.2 may be explicitly stated. By examining the asymptotic property of that constant $B(\epsilon)$ as $\epsilon \rightarrow \infty$, we prove that the minimal eigenvalue of the Weil-Petersson curvature operator on a surface of large injectivity radius is pinched by two explicit negative numbers. More precisely,

Theorem 1.3. *Let $\{X_g\}$ be a sequence of hyperbolic surfaces whose injectivity radii satisfy $\lim_{g \rightarrow \infty} \text{inj}(X_g) = \infty$. Then, for g large enough,*

$$\frac{-25}{\pi} \leq \lambda_{\min}(X_g) \leq \frac{-1}{2\pi}.$$

1.4. Bounds for small negative eigenvalues in the thick part of \mathbb{M}_g for large genus g . Thus far, we have focused exclusively on the minimal eigenvalue $\lambda_{\min}(X_g)$. In the latter portion of this part of the paper, we study some other eigenvalues of the Weil-Petersson curvature operator and their dependence on the genus g . In section 3, we restate Theorem 1.1 in [40] in terms of the eigenvalues of the Weil-Petersson curvature operator. More precisely,

Theorem 1.4. *For any $X \in \text{Teich}(S_g)$, the Weil-Petersson curvature operator \tilde{Q} at X has exactly $(3g - 3)^2$ negative eigenvalues.*

Thus, for any $X \in \text{Teich}(S_g)$, one may list the set of all non-zero eigenvalues of the Weil-Petersson curvature operator at X as follows

$$\lambda_{(3g-3)^2}(X) \leq \lambda_{(3g-3)^2-1}(X) \leq \cdots \leq \lambda_2(X) \leq \lambda_1(X) < 0.$$

The minimal eigenvalue $\lambda_{\min}(X) = \lambda_{(3g-3)^2}(X)$ basically measures the norm of the curvature operator \tilde{Q} at X .

Fixing $i \in \{1, 2, \dots, (3g - 3)^2\}$, the i -th eigenvalue $\lambda_i(X)$ is a continuous function on $\text{Teich}(S_g)$ since the Weil-Petersson metric is smooth. Thus, the Mumford compactness theorem implies that the function $\lambda_i(X)$ achieves its minima and maxima in the thick part $\text{Teich}(S_g)^{\geq \epsilon}$ for fixed $\epsilon > 0$; we denote those minima and maxima by $\underline{\lambda}_i^\epsilon(g)$ and $\bar{\lambda}_i^\epsilon(g)$ respectively. In this notation, focusing our attention now on the index i , Theorem 1.1 and Theorem 1.2 state that both $\underline{\lambda}_{(3g-3)^2}^\epsilon(g)$ and $\bar{\lambda}_{(3g-3)^2}^\epsilon(g)$ are pinched by two negative numbers which are independent of the genus of the surface. In section 6 we show that, if that index i is not close to $(3g - 3)^2$, then both $\underline{\lambda}_i^\epsilon(g)$ and $\bar{\lambda}_i^\epsilon(g)$ could be arbitrarily close to zero, once we take g large enough. More precisely,

Theorem 1.5. *If the function f satisfies $\limsup_{g \rightarrow \infty} \frac{f(g)}{9g^2} = \alpha < 1$. Then, for g sufficiently large, we have*

$$\frac{-B(\alpha, \epsilon)}{g} \leq \underline{\lambda}_{f(g)}^\epsilon(g) \leq \bar{\lambda}_{f(g)}^\epsilon(g) < 0$$

where $B(\alpha, \epsilon)$ is a constant only depending on α and ϵ . In particular,

$$\lim_{g \rightarrow \infty} \lambda_i^\epsilon(g) = \bar{\lambda}_i^\epsilon(g) = 0, \quad \text{for all } 1 \leq i \leq 8g^2.$$

A direct consequence is

Corollary 1.6. *Fix $\epsilon_0 > 0$, then for any $\epsilon > 0$, the probability*

$$\text{Prob}\{1 \leq i \leq (3g - 3)^2; |\lambda_i^{\epsilon_0}(g)| < \epsilon\} \rightarrow 1 \text{ as } g \rightarrow \infty.$$

Proof. Let $p(g) := \text{Prob}\{1 \leq i \leq (3g - 3)^2; |\lambda_i^{\epsilon_0}(g)| < \epsilon\}$. Theorem 1.5 then says that, for any $0 \leq \alpha < 1$,

$$\alpha \leq \liminf_{g \rightarrow \infty} p(g) \leq \limsup_{g \rightarrow \infty} p(g) \leq 1.$$

Since α is arbitrary in $[0, 1)$, $\lim_{g \rightarrow \infty} p(g) = 1$. \square

Mirzakhani (Theorem 4.2 in [24]) proved that for a small enough number $\epsilon_0 > 0$, the volume of the ϵ_0 -thin part of the moduli space \mathbb{M}_g is comparable to $\epsilon_0^2 \text{Vol}(\mathbb{M}_g)$ as $g \rightarrow \infty$. Corollary 1.6 suggests that the moduli space \mathbb{M}_g tends to be flat as g goes to infinity in a probabilistic sense.

1.5. Existence of holomorphic lines on sufficiently thick surfaces with uniformly pinched Weil-Petersson holomorphic sectional curvatures. In [30, 34] it was shown that the sectional curvature of the Weil-Petersson metric for $\text{Teich}(S_g)$ is negative and the holomorphic sectional curvature is bounded above by a negative number comparable to $-\frac{1}{g}$. The dependence of this bound on g begs the question as to whether there are bounds on the sectional curvature that are independent of the topology of the surface, even if one allows an additional restriction to a thick part of moduli space. However, Teo in [29] showed that, restricted to any thick part of the moduli space, the Ricci curvature is uniformly bounded from below. As the Ricci curvature is a trace over $6g - 7$ curvatures, we then see that some sectional curvatures go to zero along any sequence of surfaces X_g in the thick part of $\text{Teich}(S_g)$ as $g \rightarrow \infty$. This also suggests the question¹ whether,

Question 1. Restricted to any thick parts of the moduli spaces, do all of the sectional curvatures tend to zero as the genus goes to infinity?

The goal of the second part of this paper is to show, for the portion of the Teichmüller space composed of surfaces which are “sufficiently thick” – a term we will define precisely in Definition 2.14 – the existence of holomorphic lines whose holomorphic sectional curvatures are uniformly bounded away from zero. This gives a negative answer to the question above.

In notational preparation for the more precise version of the result, we recall that a holomorphic section of the (complexified) tangent space $T_X \text{Teich}(S_g)$

¹We are told by Zheng Huang that Question 1 is originally raised by Maryam Mirzakhani. We would like to thank both of them here.

of $X \in \text{Teich}(S_g)$ in the Teichmüller space $\text{Teich}(S_g)$ is defined by a harmonic Beltrami differential μ .

Theorem 1.7. *There is a constant ι_0 so that if $X \in \text{Teich}(S_g)$ is sufficiently thick, i.e. $\text{inj}(X) > \iota_0$, then there exists a μ_X in the tangent space $T_X \text{Teich}(S_g)$ so that the Weil-Petersson holomorphic sectional curvature $K(\mu_X)$ along μ_X satisfies*

$$K(\mu_X) \leq \frac{-81C_0}{6400 \cdot \pi^2} < 0$$

where $C_0 = \frac{2}{3C(1)^2 \cdot \text{Vol}_{\mathcal{D}}(B(0;1))}$.

The expressions $C(1)$ refers to an explicit function $C(\text{inj}(X))$ of the injectivity radius $\text{inj}(X)$ which we shall display in Definition 2.8. In addition, there is also a uniform lower bound for $K(\mu_X)$ which is due to Huang in [10] (see also Theorem 1.2).

An almost immediate corollary of our method of proof refers to a sequence of surfaces X_g of growing genus whose injectivity radii $\text{inj}(X_g) \rightarrow \infty$ as $g \rightarrow \infty$.

Corollary 1.8. *Let $\{X_g\}$ be a sequence of hyperbolic surfaces whose injectivity radii satisfy $\lim_{g \rightarrow \infty} \text{inj}(X_g) = \infty$. Then, there exists a uniform constant $E > 0$ such that for g large enough, the Weil-Petersson holomorphic sectional curvatures satisfy*

$$-\frac{2}{\pi} \leq \min_{\sigma_g \subset T_{X_g} \text{Teich}(S_g)} K(\sigma_g) \leq -E < 0$$

where the minimum runs over all the holomorphic lines in $T_{X_g} \text{Teich}(S_g)$.

[We note that the above Theorem 1.7 also proves that the minimal (most negative) eigenvalue of the Weil-Petersson curvature operator, restricted on sufficiently thick hyperbolic surface, is uniformly bounded away from zero. In that sense, this result then makes contact with, and is certainly consistent with, Theorem 1.1, but the two results are distinct, since our example here is restricted to hold on this special ‘sufficiently thick’ region (and, furthermore, the asserted constants are different).]

1.6. A remark on choice of normalizations. Some of the uniformity of the bounds that are independent of genus of course results from the conventions of fixing the uniformized metrics on the Riemann surfaces to all have curvature identically -1 (instead of, say, curvature identically $-g$), forcing the areas of the surfaces to grow linearly with g in area. Other conventions on representatives of the moduli space would translate into other bounds, but the ones we choose are the ones that seem most prevalent in the present literature.

1.7. Discussion of the methods. The methods for the two parts of the paper are distinct. In the first part, we begin with the formula of the Weil-Petersson curvature operator \tilde{Q} in [40]: see equation (3.4). That formula exhibits $\langle \tilde{Q}(A), A \rangle$ as a sum of terms involving the operator $-2(\Delta-2)^{-1}$ and algebraic expressions in a holomorphic orthonormal basis of $T_X \text{Teich}(S_g)$. By applying some fundamental properties of the operator $-2(\Delta-2)^{-1}$, we give lower and upper bounds for $\langle \tilde{Q}(A), A \rangle$ in terms of, for our applications, quantities that only involve pointwise values of the holomorphic basis. Here we briefly outline our proofs of Theorem 1.1, Theorem 1.2 and Theorem 1.7.

For the proof of **Theorem 1.1**: By applying our upper bound for the curvature operator formula to the most symmetric element $A \in \wedge^2 T_X \text{Teich}(S_g)$ we show that (up to a positive constant) $\langle \tilde{Q}(A), A \rangle$ is bounded above by the scalar curvature of the Weil-Petersson metric at X , scaled by a factor of $\frac{1}{g}$. Then we apply Wolpert's upper bound of the scalar curvature which involves a factor of g to finish the proof.

For the proof of **Theorem 1.2**: Choose an arbitrary element B of unit length in $\wedge^2 T_X \text{Teich}(S_g)$. Our lower bound of $\langle \tilde{Q}(B), B \rangle$ involves only terms that are products of our holomorphic basis, weighted by the coefficients of the element B in that basis. Well-known bounds on terms like these then yield a bound in terms of those coefficients and an explicit function of the injectivity radius. The condition that B is of unit norm then yields an expression that is independent of genus.

For the proof of **Theorem 1.7**: We choose a holomorphic line that corresponds to the image of a constant function on the disk under the Θ -operator to the hyperbolic surface X_g , here thought of as the quotient of \mathbb{H}^2 under the action of a Fuchsian group Γ_g . The resulting Poincaré series has a term corresponding to the identity element of Γ_g and a series of other 'error' terms. We adapt a method of Ahlfors [2] to show that outside a large ball, the contribution of the error terms is bounded by a subharmonic function. Taking the ball large enough, we can show that the contribution of the error terms, now estimated by its value on the boundary of the large ball – in its role as the boundary of the region of subharmonicity – can be made arbitrarily small. Thus only the term corresponding to the identity element of Γ_g cannot be made arbitrarily small, and estimates of that term yield the desired bound.

1.8. Plan of the paper. Section 2 provides some necessary background and the basic properties of the Weil-Petersson geometry of Teichmüller space that we will need. After that preparatory section, the paper splits into two parts, each of which may be read independently of the other. The first part treats our results on the Weil-Petersson curvature operator, while the second part discusses the curvature of some holomorphic sections of the tangent bundle over very thick parts of Teichmüller space.

Part I begins with section 3 in which we estimate the formula of the Weil-Petersson curvature operator and restate the non-positivity of the Weil-Petersson curvature operator in terms of eigenvalues. Section 4 provides the upper bound Theorem 1.1 for the minimal eigenvalue. In section 5 we establish Theorem 1.2, the lower bound for the minimal eigenvalue in the thick part. We prove Theorem 1.3 and Theorem 1.5 in section 6.

Part II spans two sections. We begin in section 7 with some estimates on harmonic Beltrami differentials, and then apply those estimates in section 8 to prove Theorem 1.7 and Corollary 1.8.

Acknowledgement. *The authors would like to thank Maryam Mirzakhani, Hugo Parlier and Kasra Rafi for useful conversations. They also would like to thank Scott Wolpert for useful conversations and suggestions on part (2) of this article. They deeply thank Zheng (Zeno) Huang for bringing Question 1 to their attention, his detailed comments for this article, and for his support over many years. The first author also gratefully acknowledges support from the U.S. National Science Foundation through grant DMS 1007383. This work was partially completed while the second author visited the Mathematical Sciences Center of Tsinghua university. Both authors acknowledge support from U.S. National Science Foundation grants DMS 1107452, 1107263, 1107367 “RNMS: Geometric structures And Representation varieties”(the GEAR Network).*

2. NOTATIONS AND PRELIMINARIES

2.1. Notation and Background on Weil-Petersson curvatures. In this section, we set our notations and quickly review the relevant background material on the Weil-Petersson metric and curvatures. We have two principal goals. For use in Part I, we prove Proposition 2.5, which shows that the scalar curvatures are comparable to the L^2 norms of a pointwise Bergman sum of harmonic Beltrami differentials. For use in Part II, we prove Proposition 2.11, which shows that the holomorphic sectional curvature along a harmonic Beltrami differential μ is estimated in terms of powers of its normalized L^∞ norm.

To begin, recall that we denoted by S_g a closed oriented surface of genus $g \geq 2$. We may equip S_g with a hyperbolic metric $\sigma(z)|dz|^2$, here written in a local conformal coordinate z induced by the metric. Again, $\text{Teich}(S_g)$ is the Teichmüller space of surfaces of genus g , which we may construe as equivalence classes under the action of the group Diff_0 of diffeomorphisms isotopic to the identity of the space of hyperbolic surfaces $X = (S_g, \sigma|dz|^2)$. The tangent space $T_X \text{Teich}(S_g)$ at a point $X = (S_g, \sigma|dz|^2)$ is identified with the space of *harmonic Beltrami differentials* on X , i.e. forms on X expressible as $\mu = \frac{\psi}{\sigma}$ where $\psi \in H^0(X, K^2)$ is a holomorphic quadratic differential on X . Let $z = x + iy$ and $dA = \sigma(z)dxdy$ be the volume form. The *Weil-Petersson metric* is the Hermitian metric on $\text{Teich}(S_g)$ arising

from the the *Petersson scalar product*

$$\langle \varphi, \psi \rangle = \int_S \frac{\varphi \cdot \bar{\psi}}{\sigma^2} dA$$

via duality. We will concern ourselves primarily with its Riemannian part g_{WP} . Throughout this paper we denote the Teichmüller space endowed with the Weil-Petersson metric by $\text{Teich}(S_g)$.

Set $D = -2(\Delta - 2)^{-1}$ where Δ is the Beltrami-Laplace operator on $X = (S, \sigma|dz|^2)$. The operator D is positive and self-adjoint. The following inequality follows from the maximum principle; see Lemma 5.1 in [32] for details.

Lemma 2.1 ([32]). *For any harmonic Beltrami differential μ on X , we have*

$$D(|\mu|^2) \geq \frac{|\mu|^2}{3}.$$

The following property is well-known to experts – see for example Lemma 4.3 in [16]. For completeness, we include the proof.

Lemma 2.2. *Let D be the operator above. Then, for any complex-valued function $f \in C^\infty(X)$,*

$$0 \leq \int_X (D(f)\bar{f})dA \leq \int_X |f|^2 dA.$$

Proof. Let $g = D(f)$, so that $f = \frac{-1}{2}(\Delta - 2)g$. The left inequality follows directly by integrating by parts.

For the right inequality, begin by assuming that f is real-valued. In that case decompose $f = \sum_{i=0}^{\infty} \phi_i$ as a linear combination of eigenfunctions of the Laplacian: here ϕ_i satisfies $\int_X (\phi_i \phi_j) dA = 0$ for all $i \neq j$ and $\Delta \phi_i = \mu_i \phi_i$, where $\mu_i < 0$ is the i -th eigenvalue of the Beltrami-Laplace operator Δ . Since $\int_X (\phi_i \phi_j) dA = 0$ for all $i \neq j$, we have

$$\begin{aligned} \int_X (D(f)f)dA &= \int_X \left(\sum_{i=0}^{\infty} \frac{2\phi_i}{2 - \mu_i} \right) \left(\sum_{i=0}^{\infty} \phi_i \right) dA \\ &= \int_X \sum_{i=0}^{\infty} \frac{2}{2 - \mu_i} |\phi_i|^2 dA \\ &\leq \int_X \sum_{i=0}^{\infty} |\phi_i|^2 dA \\ &= \int_X f^2 dA. \end{aligned}$$

If f is complex-valued, one applies the same method to the real and imaginary parts of f separately, along with a standard use of the self-adjointness of D . \square

We conclude this discussion of the operator $D = -2(\Delta - 2)^{-1}$ by recording some elementary properties of its Green's function.

Proposition 2.3. *Let D be the operator $D = -2(\Delta - 2)^{-1}$. Then there exists a Green function $G(w, z)$ for D satisfying:*

- (1). $D(f)(z) = \int_{w \in X} G(z, w) f(w) dA(w)$ for any $f \in C^\infty((X, \sigma|dz|^2), \mathbb{C})$.
- (2). $G(w, z)$ is positive.
- (3). $G(w, z)$ is symmetric, i.e., $G(w, z) = G(z, w)$.

Proof. See for example [26] and [34]. □

2.2. The Riemannian tensor of the Weil-Petersson metric. The curvature tensor of the Weil-Petersson metric is given as follows. As described in the opening paragraph of section 2.1, let μ_i, μ_j be two elements in the tangent space $T_X \text{Teich}(S_g)$ at X , so that the metric tensor might be written in local coordinates as

$$g_{i\bar{j}} = \int_X \mu_i \cdot \bar{\mu}_j dA.$$

For the inverse of $(g_{i\bar{j}})$, we use the convention

$$g^{i\bar{j}} g_{k\bar{j}} = \delta_{ik}.$$

Then the curvature tensor is given by

$$R_{i\bar{j}k\bar{l}} = \frac{\partial^2}{\partial t^k \partial \bar{t}^l} g_{i\bar{j}} - g^{s\bar{t}} \frac{\partial}{\partial t^k} g_{i\bar{s}} \frac{\partial}{\partial \bar{t}^l} g_{s\bar{j}}.$$

The following curvature formula was established in [30, 34]. One can also see [13] for a derivation from a third perspective. It has been applied to study various curvature properties of the Weil-Petersson metric. In [27] Schumacher showed that $\text{Teich}(S_g)$ has strongly negative curvature in the sense of Siu. Huang in his thesis [9] showed that the sectional curvatures of $\text{Teich}(S_g)$ can not be bounded away from zero. Liu-Sun-Yau in [15] showed that $\text{Teich}(S_g)$ has dual Nakano negative curvature, which says that the complex curvature operator on the dual tangent bundle is positive in some sense. Motivated by the method in [15], the second author in [40] showed that the $\text{Teich}(S_g)$ has negative semi-definite Riemannian curvature operator. One can also see [16, 17, 18, 29, 37, 39] for other aspects of the curvature of $\text{Teich}(S_g)$.

Theorem 2.4 (Tromba, Wolpert). *The curvature tensor satisfies*

$$R_{i\bar{j}k\bar{l}} = \int_X D(\mu_i \mu_{\bar{j}}) \cdot (\mu_k \mu_{\bar{l}}) dA + \int_X D(\mu_i \mu_{\bar{l}}) \cdot (\mu_k \mu_{\bar{j}}) dA.$$

2.2.1. *Weil-Petersson Scalar Curvature.* Let $\{\mu_i\}_{i=1}^{3g-3}$ be a holomorphic orthonormal basis representing the tangent space $T_X \text{Teich}(S_g)$ at X . Then the Ricci curvature of $\text{Teich}(S_g)$ at X in the direction μ_i is given by

$$\begin{aligned} Ric(\mu_i) &= - \sum_{j=1}^{3g-3} R_{i\bar{j}j\bar{i}} \\ &= - \sum_{j=1}^{3g-3} \left(\int_X D(\mu_i \mu_{\bar{j}}) \cdot (\mu_j \mu_{\bar{i}}) dA + \int_X D(|\mu_i|^2) \cdot (|\mu_j|^2) dA \right). \end{aligned}$$

Since the scalar curvature $Sca(X)$ at X is the trace of the Ricci tensor, we find we may express the scalar curvature as

$$(2.1) \quad Sca(X) = - \sum_{i=1}^{3g-3} \sum_{j=1}^{3g-3} \left(\int_X D(\mu_i \mu_{\bar{j}}) \cdot (\mu_j \mu_{\bar{i}}) dA + \int_X D(|\mu_i|^2) \cdot (|\mu_j|^2) dA \right).$$

From this expression for the scalar curvature, we may apply our estimates for the operator D in section 2.1 to obtain estimates for the scalar curvature from above and below in terms of pointwise-defined quantities.

Proposition 2.5. *For any $X \in \text{Teich}(S_g)$, the scalar curvature $Sca(X)$ at X satisfies*

$$-2 \int_X \left(\sum_{i=1}^{3g-3} |\mu_i|^2 \right)^2 dA \leq Sca(X) \leq -\frac{1}{3} \int_X \left(\sum_{i=1}^{3g-3} |\mu_i|^2 \right)^2 dA$$

where $\{\mu_i\}_{i=1}^{3g-3}$ is any holomorphic orthonormal basis of the tangent space at X .

Proof. We begin with the right-hand-side inequality. By Lemma 2.1, the first term in the expression (2.1) has a sign, so that we can then apply Lemma 2.2 to find

$$\begin{aligned} -Sca(X) &= \sum_{i=1}^{3g-3} \sum_{j=1}^{3g-3} \left(\int_X D(\mu_i \mu_{\bar{j}}) \cdot (\mu_j \mu_{\bar{i}}) dA + \int_X D(|\mu_i|^2) \cdot (|\mu_j|^2) dA \right) \\ &\geq \sum_{i=1}^{3g-3} \sum_{j=1}^{3g-3} \int_X D(|\mu_i|^2) \cdot (|\mu_j|^2) dA \\ &\geq \frac{1}{3} \int_X \left(\sum_{i=1}^{3g-3} |\mu_i|^2 \right)^2 dA. \end{aligned}$$

For the left hand side inequality, we use the right side of the estimate in Lemma 2.2 twice, with the preliminary step of bringing the sum into the

integrands. We compute

$$\begin{aligned}
-Sca(X) &= \sum_{i=1}^{3g-3} \sum_{j=1}^{3g-3} \left(\int_X D(\mu_i \mu_{\bar{j}}) \cdot (\mu_j \mu_{\bar{i}}) dA + \int_X D(|\mu_i|^2) \cdot (|\mu_j|^2) dA \right) \\
&= \sum_{i=1}^{3g-3} \sum_{j=1}^{3g-3} \int_X D(\mu_i \mu_{\bar{j}}) \cdot (\mu_j \mu_{\bar{i}}) dA + \int_X \sum_{i=1}^{3g-3} \sum_{j=1}^{3g-3} D(|\mu_i|^2) \cdot (|\mu_j|^2) dA \\
&\leq \sum_{i=1}^{3g-3} \sum_{j=1}^{3g-3} \int_X (\mu_i \mu_{\bar{j}}) \cdot (\mu_j \mu_{\bar{i}}) dA + \int_X D\left(\sum_{i=1}^{3g-3} |\mu_i|^2\right) \cdot \left(\sum_{j=1}^{3g-3} |\mu_j|^2\right) dA \\
&= \sum_{i=1}^{3g-3} \sum_{j=1}^{3g-3} \int_X |\mu_i|^2 |\mu_j|^2 dA + \int_X D\left(\sum_{i=1}^{3g-3} |\mu_i|^2\right) \cdot \left(\sum_{i=1}^{3g-3} |\mu_i|^2\right) dA \\
&\leq 2 \int_X \left(\sum_{i=1}^{3g-3} |\mu_i|^2\right)^2 dA.
\end{aligned}$$

□

Remark 2.6. The Cauchy-Schwarz inequality applied to the right hand side estimate in Proposition 2.5 leads to a numerical lower bound for $-Sca(X)$ in the following way:

$$\begin{aligned}
-Sca(X) &\geq \frac{\int_X \left(\sum_{i=1}^{3g-3} |\mu_i|^2\right)^2 dA}{3} \\
&\geq \frac{\left(\int_X \left(\sum_{i=1}^{3g-3} |\mu_i|^2\right) dA\right)^2}{3 \text{Area}(S)} = \frac{3(g-1)}{4\pi}.
\end{aligned}$$

Now, without using Lemma 2.1, Wolpert in [34] proved a better lower bound as $-Sca(X) \geq \frac{3(3g-2)}{4\pi}$ by expanding $\sum_{i=1}^{3g-3} |\mu_i|^2$ (and $D(\sum_{i=1}^{3g-3} |\mu_i|^2)$) in the definition (2.1) in terms of eigenfunctions of Δ at X . On the other hand, our goal in Proposition 2.5 is to estimate $-Sca(X)$ is estimated by the integral $\int_X \left(\sum_{i=1}^{3g-3} |\mu_i|^2\right)^2 dA$, an expression we will apply later, so we content ourselves with the bound above.

2.2.2. Weil-Petersson holomorphic sectional curvatures. Recall the holomorphic sectional curvature is a sectional curvature along a holomorphic line. Let $\mu \in T_X \text{Teich}(S_g)$. Then Theorem 2.4 tells that the holomorphic sectional curvature $K(\mu)$ along the holomorphic line spanned by μ is

$$K(\mu) = \frac{-2 \cdot \int_X D(|\mu|^2) \cdot (|\mu|^2) dA}{\|\mu\|_{WP}^4}.$$

From Lemma 2.1, Lemma 2.2 and the equation above we have

Proposition 2.7. *For any $\mu \in T_X \text{Teich}(S_g)$, the holomorphic sectional curvature $K(\mu)$ satisfies*

$$-\frac{2 \cdot \int_X |\mu|^4 dA}{\|\mu\|_{WP}^4} \leq K(\mu) \leq -\frac{2 \cdot \int_X |\mu|^4 dA}{3\|\mu\|_{WP}^4}.$$

We will refer several times to the a constant $C(\text{inj})$ depending on the injectivity radius, which we pause now to define.

Definition 2.8. Set $C(\text{inj}(X)) = \left(\frac{4\pi}{3} \left(1 - \left(\frac{4e^{\text{inj}(X)}}{(1+e^{\text{inj}(X)})^2}\right)^3\right)\right)^{-1}$.

Remark 2.9. We note here that as the injectivity radius $\text{inj}(X)$ tends to zero, the constant $C(\text{inj}(X)) = \frac{1}{\pi \text{inj}(X)^2} + o\left(\frac{1}{\text{inj}(X)^2}\right)$. And $C(\text{inj}(X))$ tends to $\frac{3}{4\pi}$ as $\text{inj}(X)$ goes to infinity.

Next we recall the following proposition which is implicitly proved in [10, 29, 39]. For the sake of keeping the exposition self-contained, we give the outline of the proof which follows the identical argument as that of Proposition 3.1 in [29].

Proposition 2.10. *Let $(X, \sigma|dz|^2)$ be a closed hyperbolic surface and $\mu \in T_X \text{Teich}(S_g)$ be a harmonic Beltrami differential of X . Then, for any $q \in X$ we have*

$$|\mu(q)|^2 \leq C(r) \int_{B(q;r)} |\mu(z)|^2 dA(z) \quad \forall 0 < r \leq \text{inj}(q).$$

Where $\text{inj}(q)$ is the injectivity radius of q in X and the explicit constant $C(r)$ is given in Definition 2.8.

Proof. Let $\mu \in T_X \text{Teich}(S_g)$; we denote its lift into the hyperbolic disk by ν . For any point $q \in X$, we may conjugate the lift by a linear fractional transformation so that

$$\nu(0) = \mu(q).$$

For any $r \in (0, \text{inj}(q)]$, elementary hyperbolic geometry gives that the hyperbolic disk of radius r centered at 0 is

$$B(0; r) = \left\{z \in \mathcal{D}; |z| \leq \frac{e^r - 1}{e^r + 1}\right\}.$$

Since ν is a harmonic Beltrami differential on the disk, there exists a holomorphic function f such that

$$\nu(z) = \frac{\overline{f(z)}}{\frac{4}{(1-|z|^2)^2}} = \frac{\overline{f(z)}(1-|z|^2)^2}{4}.$$

Of course, since f is holomorphic, we may expand it as

$$f(z) = \sum_{n=0}^{\infty} a_n z^n$$

where a_n are complex numbers. In particular we have

$$|\mu(q)| = |\nu(0)| = \frac{|a_0|}{4}.$$

Therefore, for any $r \in (0, \text{inj}(q)]$, we have

$$\begin{aligned} \int_{B(q;r)} |\mu(z)|^2 dA(z) &= \int_{B(0;r)} |\nu(z)|^2 dA(z) \\ &= \int_{B(0;r)} |f(z)|^2 \frac{(1-|z|^2)^2}{4} |dz|^2 \\ &= \int_0^{\frac{e^r-1}{e^r+1}} \int_0^{2\pi} \left| \sum_{n=0}^{\infty} a_n u^n e^{in\theta} \right|^2 \frac{(1-u^2)^2}{4} u du d\theta \\ &= \frac{\pi}{2} \int_0^{\frac{e^r-1}{e^r+1}} \sum_{n=0}^{\infty} |a_n|^2 u^{2n} (1-u^2)^2 u du \\ &\geq \frac{\pi}{2} |a_0|^2 \int_0^{\frac{e^r-1}{e^r+1}} (1-u^2)^2 u du \\ &= \frac{4\pi}{3} \left(1 - \left(\frac{4e^r}{(1+e^r)^2}\right)^3\right) |\mu(q)|^2. \end{aligned}$$

The conclusion follows by dividing the constant on both sides. \square

We combine these last two propositions to obtain the following proposition which is crucial in the proof of Theorem 1.7 (which produces a holomorphic section whose curvature is pinched): this proposition traps the holomorphic sectional curvature between powers of L^∞ bounds on (normalized) holomorphic Beltrami differentials.

Proposition 2.11. *Let $X \in \text{Teich}(S_g)$ with $\text{inj}(X) \geq 1$. Then for any $\mu \in T_X \text{Teich}(S_g)$, there exists a constant $C_0 > 0$, independent of the genus, such that the holomorphic sectional curvature $K(\mu)$ satisfies*

$$-\frac{2 \cdot \sup_{z \in X} |\mu(z)|^2}{\|\mu\|_{WP}^2} \leq K(\mu) \leq -\frac{C_0 \cdot \sup_{z \in X} |\mu(z)|^4}{\|\mu\|_{WP}^4}.$$

Proof. We begin with the left hand side. By Proposition 2.7 we have

$$-K(\mu) \leq \frac{2 \cdot \sup_{z \in X} |\mu(z)|^2 \cdot \int_X |\mu|^2 dA}{\|\mu\|_{WP}^4} = \frac{2 \cdot \sup_{z \in X} |\mu(z)|^2}{\|\mu\|_{WP}^2}.$$

For the right hand side, since X is compact, we let $p \in X$ such that $|\mu(p)| = \sup_{z \in X} |\mu(z)|$. Let $B(p; 1)$ be the closed ball of hyperbolic radius 1 centered at p . Use the Cauchy-Schwarz inequality to obtain

$$\int_X |\mu|^4 dA \geq \int_{B(p;1)} (|\mu|^4) dA \geq \frac{(\int_{B(p;1)} |\mu|^2 dA)^2}{\text{Vol}(B(p; 1))}.$$

Applying Proposition 2.10 to the last term, we find

$$\int_X |\mu|^4 dA \geq \frac{|\mu(p)|^4}{C(1)^2 \cdot \text{Vol}(B(p; 1))} = \frac{\sup_{z \in X} |\mu(z)|^4}{C(1)^2 \cdot \text{Vol}(B(p; 1))}.$$

Finally, by choosing $C_0 = \frac{2}{3C(1)^2 \cdot \text{Vol}(B(p; 1))}$ (where $C(1)$ is the explicit constant from Definition 2.8), the conclusion follows from Proposition 2.7. \square

Remark 2.12. Combining Proposition 2.10 and Proposition 2.11, one may conclude that the holomorphic sectional curvatures of the ϵ_0 -thick part of the moduli space are uniformly bounded below by a constant only depending on ϵ_0 , which was proved in [10].

Remark 2.13. If one would like to search for harmonic Beltrami differentials with uniformly negative holomorphic sectional curvatures as the genus goes to infinity, Proposition 2.11 tells that it suffices to find such differentials with Weil-Petersson L^2 norms bounded above and Weil-Petersson L^∞ norms bounded below. We develop this theme in the second part of this paper.

2.3. Sufficiently thick surfaces. Often in Teichmüller theory, one studies problems in the setting where some curve α on the surface S has hyperbolic X -length $\ell_X(\alpha)$ small, say $\ell_X(\alpha) \leq \epsilon_0$. Alternatively, also often studies issues where the surface has injectivity radius $\text{inj}(X)$ bounded away from zero, say $\text{inj}(X) \geq \epsilon_1$. In this paper, a number of our results are in the region where the hyperbolic surface X has a large injectivity radius, where 'large' here means large enough for some *a priori* bounds to apply.

Definition 2.14. For a given large constant $C > 0$, a surface X is *C -sufficiently thick* or *sufficiently thick* (if the constant C is clear from the context) if $\text{inj}(X) \geq C$.

We remark that there often is a non-empty subspace $\mathbb{M}^{\geq C}$ of sufficiently thick surfaces of the moduli space \mathbb{M} . It is well-known that there exist sequences of hyperbolic surfaces whose injectivity radii grow without bounds. For example, the fundamental group of a closed hyperbolic surface X is residually finite (one can see [8] for details). After taking finite normal covers to remove simple closed curves of bounded lengths in X , we can find a sequence of hyperbolic surfaces X_{g_k} with injectivity radii $\text{inj}(X_{g_k}) \rightarrow \infty$ as $g_k \rightarrow \infty$. Moreover, Buser and Sarnak [4] proved that there exists a family of closed surfaces X_k of genus g_k with $g_k \rightarrow \infty$ as $k \rightarrow \infty$ and $\text{inj}(X_k) \geq \frac{2 \ln g_k}{3}$.

Part 1. Bounds on the Weil-Petersson curvature operator

3. THE WEIL-PETERSSON CURVATURE OPERATOR AT X AND ITS EIGENVALUES

In this section, we begin our study of the Weil-Petersson curvature operator \tilde{Q} , establishing some preliminary upper and lower bounds for $<$

$\tilde{Q}(A), A >$ in Proposition 3.6. Those estimates rely on a formula for $< \tilde{Q}(A), A >$ displayed in Proposition 3.5 which is particularly well-adapted for estimations. The first subsection provides notation and context.

Before we study the curvature operator on $\text{Teich}(S_g)$, we set some notation. Let $U \subset \text{Teich}(S_g)$ be a neighborhood of X in Teichmüller space, and let $(t_1, t_2, \dots, t_{3g-3})$ be a system of local holomorphic coordinates on U such that $\{t_i(X) = \mu_i\}_{1 \leq i \leq 3g-3}$ is holomorphic orthonormal at X with $t_i = x_i + \mathbf{i}y_i$ ($1 \leq i \leq 3g-3$). Then $(x_1, x_2, \dots, x_{3g-3}, y_1, y_2, \dots, y_{3g-3})$ are real smooth coordinates in U which relate to the complex coordinates as

$$\frac{\partial}{\partial x_i} = \frac{\partial}{\partial t_i} + \frac{\partial}{\partial \bar{t}_i}, \quad \frac{\partial}{\partial y_i} = \mathbf{i} \left(\frac{\partial}{\partial t_i} - \frac{\partial}{\partial \bar{t}_i} \right).$$

Let $T \text{Teich}(S_g)$ be the real tangent bundle of $\text{Teich}(S_g)$ and $\wedge^2 T \text{Teich}(S_g)$ be the exterior wedge product of $T \text{Teich}(S_g)$ with itself. For any $X \in U$, we have

$$T_X \text{Teich}(S_g) = \text{Span} \left\{ \frac{\partial}{\partial x_i}(X), \frac{\partial}{\partial y_j}(X) \right\}_{1 \leq i, j \leq 3g-3}.$$

and

$$\wedge^2 T \text{Teich}(S_g) = \text{Span} \left\{ \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j}, \frac{\partial}{\partial x_k} \wedge \frac{\partial}{\partial y_l}, \frac{\partial}{\partial y_m} \wedge \frac{\partial}{\partial y_n} \right\}.$$

The space $\wedge^2 T \text{Teich}(S_g)$ has (real) dimension $(3g-3)(6g-7)$. Let $<, >$ refer to the pairing of vectors with respect to the Weil-Petersson metric. The natural inner product on $\wedge^2 T \text{Teich}(S_g)$, associated to the Weil-Petersson metric, is given by

$$(3.1) \quad \langle V_1 \wedge V_2, V_3 \wedge V_4 \rangle_{eu} := \langle V_1, V_3 \rangle \langle V_2, V_4 \rangle - \langle V_1, V_4 \rangle \langle V_2, V_3 \rangle$$

and extended linearly, where V_i are real vectors. One may refer to ([11], p. 238) for more details.

In terms of these real coordinates, the Weil-Petersson curvature operator \tilde{Q} may be described in the following way. Let $X \in \text{Teich}(S_g)$, let R be the Riemannian curvature tensor of the Weil-Petersson metric, and let $\{e_i\}_{i=1}^{6g-6}$ be an orthonormal basis of $T_X \text{Teich}(S_g)$. Set

$$R_{ijkl} := \langle R(e_i, e_j)e_k, e_l \rangle.$$

It is clear that

$$\wedge^2 T_X \text{Teich}(S_g) = \text{Span} \{e_i \wedge e_j\}_{1 \leq i < j \leq (6g-6)}.$$

Then the operator $\tilde{Q} : \wedge^2 T_X \text{Teich}(S_g) \rightarrow \wedge^2 T_X \text{Teich}(S_g)$ may be written in this notation as

$$\tilde{Q} \left(\sum_{1 \leq i < j \leq (6g-6)} a_{ij} e_i \wedge e_j \right) := \sum_{1 \leq i < j \leq (6g-6)} \sum_{1 \leq k < l \leq (6g-6)} a_{ij} R_{ijkl} e_k \wedge e_l,$$

where the coefficients a_{ij} are set to be real.

From equation (3.1) it is easy to see that $\{e_i \wedge e_j\}_{1 \leq i < j \leq (6g-6)}$ is an orthonormal basis of $\wedge^2 T_X \text{Teich}(S_g)$. Then,

$$\begin{aligned} & \langle \tilde{Q}(\sum_{1 \leq i < j \leq (6g-6)} a_{ij} e_i \wedge e_j), \sum_{1 \leq i < j \leq (6g-6)} b_{ij} e_i \wedge e_j \rangle_{eu} \\ &= \sum_{1 \leq i < j \leq (6g-6)} \sum_{1 \leq k < l \leq (6g-6)} a_{ij} b_{kl} R_{ijkl}, \end{aligned}$$

where here again the coefficients b_{ij} are real.

We define the *associated (bilinear, symmetric) curvature form* to be the bilinear form Q on $\wedge^2 T \text{Teich}(S_g)$ given by $Q(V_1 \wedge V_2, V_3 \wedge V_4) = R(V_1, V_2, V_3, V_4)$ and extended linearly, where the V_i are real vectors. It is easy to see that Q is a bilinear symmetric form (one can see more details in [14]).

In this notation, we see that we may write $Q(A, A)$ as $Q(A, A) = \langle \tilde{Q}(A), A \rangle_{eu}$ for all $A \in \wedge^2 T \text{Teich}(S_g)$. By the symmetry of the Riemannian curvature tensor and the definition of the scalar curvature, we then find

Lemma 3.1. (1). \tilde{Q} is self-adjoint.

(2). The trace $\text{Tr}(\tilde{Q})(X)$ of \tilde{Q} at X satisfies $\text{Tr}(\tilde{Q})(X) = \text{Sca}(X)$.

For more details on the Riemannian curvature operator, one can see section 2.2 in [25].

3.1. The eigenvalues of the Weil-Petersson curvature operator. The action of the almost complex structure \mathbf{J} on $T_X \text{Teich}(S_g)$ extends to a natural action of \mathbf{J} on $\wedge^2 T_X \text{Teich}(S_g)$, defined as follows on a basis

$$\begin{cases} \mathbf{J} \circ \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j} := \frac{\partial}{\partial y_i} \wedge \frac{\partial}{\partial y_j}, \\ \mathbf{J} \circ \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j} := -\frac{\partial}{\partial y_i} \wedge \frac{\partial}{\partial x_j} = \frac{\partial}{\partial x_j} \wedge \frac{\partial}{\partial y_i}, \\ \mathbf{J} \circ \frac{\partial}{\partial y_i} \wedge \frac{\partial}{\partial y_j} := \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j}, \end{cases}$$

and then extended linearly. It is easy to see that, as an operator on $\wedge^2 T_X \text{Teich}(S_g)$, we have $\mathbf{J} \circ \mathbf{J} = id$.

Part of the second author's thesis shows that for any $X \in \text{Teich}(S_g)$, the curvature operator \tilde{Q} at X is negative semi-definite. More precisely,

Theorem 3.2 ([40]). *Let $S = S_g$ be a closed surface of genus $g > 1$ and $\text{Teich}(S_g)$ be the Teichmüller space of S endowed with the Weil-Petersson metric. And let \mathbf{J} be the almost complex structure on $\text{Teich}(S_g)$ and Q be the associated curvature form on $\text{Teich}(S_g)$. Then, for any $X \in \text{Teich}(S_g)$, we have*

(1). \tilde{Q} is negative semi-definite, i.e., $\langle \tilde{Q}(A), A \rangle_{eu} = Q(A, A) \leq 0$ for all $A \in \wedge^2 T_X \text{Teich}(S_g)$.

(2). $Q(A, A) = 0$ if and only if there exists an element B in $\wedge^2 T_X \text{Teich}(S_g)$ such that $A = B - \mathbf{J} \circ B$.

Recall that λ is called an *eigenvalue* of \tilde{Q} if there exists an element $A \in \wedge^2 T_X \text{Teich}(S_g)$ such that $\tilde{Q}(A) = \lambda \cdot A$. In that case, the element A is called

the eigenvector associated to λ . Since \tilde{Q} is self-adjoint, all of the eigenvalues of \tilde{Q} are real. Since $\dim(\text{Teich}(S_g)) = 6g - 6$, we see from Theorem 3.2 that there exist $(3g - 3)(6g - 7)$ non-positive eigenvalues $\{\nu_i\}_{i=1}^{(3g-3)(6g-7)}$ of \tilde{Q} where we set $\nu_{i+1} \leq \nu_i \leq 0$ for all $0 \leq i \leq ((3g - 3)(6g - 7) - 1)$. Since the Weil-Petersson sectional curvature is negative (see [34]), not all $\{\nu_i\}$ vanish. Our focus is on the non-zero eigenvalues. Refining the analysis of Theorem 3.2, we prove that the number of non-zero eigenvalues of the Weil-Petersson curvature operator \tilde{Q} on Teichmüller space $\text{Teich}(S_g)$ is constant.

Theorem 3.3. *For any $X \in \text{Teich}(S_g)$, the Weil-Petersson curvature operator \tilde{Q} at X has exactly $(3g - 3)^2$ negative eigenvalues.*

To prepare for the proof of Theorem 3.3, we note the following standard lemma.

Lemma 3.4. *For any $B \in \wedge^2 T_X \text{Teich}(S_g)$, the element A defined by $A = B - \mathbf{J} \circ B$ is a 0-eigenvector, i.e. $\tilde{Q}(A) = 0$.*

Proof. Of course, by part (2) of Theorem 3.2 we have $Q(A, A) = 0$. Now let $C \in \wedge^2 T_X \text{Teich}(S_g)$. By part (1) of Theorem 3.2, for all $t \in \mathbb{R}$, we have

$$\begin{aligned} 0 &\geq Q(C + tA, C + tA) = \langle Q(C + tA), C + tA \rangle_{eu} \\ &= \langle \tilde{Q}(C), C \rangle_{eu} + 2t \langle \tilde{Q}(A), C \rangle_{eu}. \end{aligned}$$

Since t is arbitrary, we see that we could choose t to make the expression $\langle \tilde{Q}(C), C \rangle_{eu} + 2 \langle \tilde{Q}(A), C \rangle_{eu} t$ above positive unless $\langle \tilde{Q}(A), C \rangle_{eu} = 0$. The conclusion then follows by choosing $C = \tilde{Q}(A)$ so that we have $0 = \langle \tilde{Q}(A), C \rangle_{eu} = \langle \tilde{Q}(A), \tilde{Q}(A) \rangle_{eu}$, yielding $\tilde{Q}(A) = 0$. \square

Proof of Theorem 3.3. It suffices to show that the dimension of the space of zero-eigenvectors of \tilde{Q} is equal to $(3g - 3)(3g - 4)$. First, it is clear that the elements in $\{\frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j} - \frac{\partial}{\partial y_i} \wedge \frac{\partial}{\partial y_j}\}_{1 \leq i < j \leq (3g-3)}$, $\{\frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j} - \frac{\partial}{\partial x_j} \wedge \frac{\partial}{\partial y_i}\}_{1 \leq i < j \leq (3g-3)}$ are linearly independent in $\wedge^2 T_X \text{Teich}(S_g)$. By Lemma 3.4, we see that $\text{Span}\{\{\frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j} - \frac{\partial}{\partial y_i} \wedge \frac{\partial}{\partial y_j}\}_{1 \leq i < j \leq (3g-3)}, \{\frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j} - \frac{\partial}{\partial x_j} \wedge \frac{\partial}{\partial y_i}\}_{1 \leq i < j \leq (3g-3)}\}$ is contained in the space of zero-eigenvectors of \tilde{Q} . Thus, $\dim\{A \in \wedge^2 T_X \text{Teich}(S_g); \tilde{Q}(A) = 0\} \geq 2 \cdot (3g-3)(3g-4)/2 = (3g-3)(3g-4)$.

On the other hand, let $A = \sum_{ij} (a_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j} + b_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j} + c_{ij} \frac{\partial}{\partial y_i} \wedge \frac{\partial}{\partial y_j}) \in \wedge^2 T_X \text{Teich}(S_g)$ with $\tilde{Q}(A) = 0$. By part (2) of Theorem 3.2, we know that $A = B - \mathbf{J} \circ B$ for some $B \in \wedge^2 T_X \text{Teich}(S_g)$. Then we have, for all $1 \leq i, j \leq (3g - 3)$,

$$a_{ij} + c_{ij} = a_{ji} + c_{ji}, \quad b_{ij} + b_{ji} = 0.$$

We rewrite it as

$$a_{ij} - a_{ji} = -(c_{ij} - c_{ji}), \quad b_{ij} = -b_{ji}.$$

Thus,

$$A = \sum_{i < j} (a_{ij} - a_{ji}) \left(\frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j} - \frac{\partial}{\partial y_i} \wedge \frac{\partial}{\partial y_j} \right) + \sum_{i < j} b_{ij} \left(\frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j} - \frac{\partial}{\partial x_j} \wedge \frac{\partial}{\partial y_i} \right).$$

In particular $A \in \text{Span}\left\{ \left\{ \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j} - \frac{\partial}{\partial y_i} \wedge \frac{\partial}{\partial y_j} \right\}_{1 \leq i < j \leq (3g-3)}, \left\{ \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j} - \frac{\partial}{\partial x_j} \wedge \frac{\partial}{\partial y_i} \right\}_{1 \leq i < j \leq (3g-3)} \right\}$. Since A is arbitrary with $\tilde{Q}(A) = 0$,

$$\dim\{A \in \wedge^2 T_X \text{Teich}(S_g); \tilde{Q}(A) = 0\} \leq 2 \cdot (3g-3)(3g-4)/2 = (3g-3)(3g-4).$$

Therefore,

$$\dim\{A \in \wedge^2 T_X \text{Teich}(S_g); \tilde{Q}(A) = 0\} = (3g-3)(3g-4).$$

□

From Theorem 3.2 and Theorem 3.3, we know that for any $X \in \text{Teich}(S_g)$, the Weil-Petersson curvature operator \tilde{Q} at X has $(3g-3)^2$ negative eigenvalues. We denote them by

$$\lambda_{(3g-3)^2}(X) \leq \lambda_{(3g-3)^2-1}(X) \leq \lambda_{(3g-3)^2-2}(X) \leq \cdots \leq \lambda_2(X) \leq \lambda_1(X) < 0$$

We close this subsection by rewriting the smallest eigenvalue $\lambda_{(3g-3)^2}(X)$ as follows

$$(3.2) \quad \lambda_{\min}(X) := \lambda_{(3g-3)^2}(X) = \min_{A \in \wedge^2 T_X \text{Teich}(S_g), \|A\|_{eu}=1} Q(A, A).$$

3.2. The Weil-Petersson curvature operator formula. In this section, we recall a formula from [40] for the curvature operator applied to an element $A \in \wedge^2 T_X \text{Teich}(S_g)$ of $\wedge^2 T_X \text{Teich}(S_g)$. This formula is particularly well-suited to estimating the associated curvature form $Q(A, A)$, and so from the formula we derive some estimates that we will need in the later sections.

To begin, we express an arbitrary element $A \in \wedge^2 T_X \text{Teich}(S_g)$ in coordinates as

$$(3.3) \quad A = \sum_{ij} (a_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j} + b_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j} + c_{ij} \frac{\partial}{\partial y_i} \wedge \frac{\partial}{\partial y_j}) \in \wedge^2 T_X \text{Teich}(S_g)$$

where a_{ij}, b_{ij}, c_{ij} are real numbers. The following formula is crucial not only for our analysis in this section, but indeed for much of our work in this part of the paper.

Proposition 3.5. *With A defined as above, we may write $Q(A, A)$ as*

$$(3.4) \quad \begin{aligned} Q(A, A) = & -4 \int_X D(\text{Im}\{F(z, z) + \mathbf{i}H(z, z)\}) \cdot (\text{Im}\{F(z, z) + \mathbf{i}H(z, z)\}) dA(z) \\ & - 2 \int_{X \times X} G(z, w) |F(z, w) + \mathbf{i}H(z, w)|^2 dA(w) dA(z) \\ & + 2 \text{Re} \left\{ \int_{X \times X} G(z, w) (F(z, w) + \mathbf{i}H(z, w))(F(w, z) + \mathbf{i}H(w, z)) dA(w) dA(z) \right\} \end{aligned}$$

where $G(z, w)$ is the Green's function for the operator D , the expression $F(z, w)$ is set to be $F(z, w) = \sum_{i,j=1}^{3g-3} (a_{ij} + c_{ij}) \mu_i(w) \cdot \overline{\mu_j(z)}$ and $H(z, w)$ is defined as $H(z, w) = \sum_{i,j=1}^{3g-3} b_{ij} \mu_i(w) \cdot \overline{\mu_j(z)}$.

Proof. The conclusion directly follows from Proposition 4.1 and Proposition 4.3 in [40]. \square

Using this decomposition of $Q(A, A)$, we estimate its value in the following proposition. The proof uses a similar idea as in the proof of Theorem 4.4 in [40].

Proposition 3.6. *Under the same conditions as in Proposition 3.5, we have*

- (1). $-Q(A, A) \geq 4 \int_X D(\operatorname{Im}\{F(z, z) + \mathbf{i}H(z, z)\}) \cdot (\operatorname{Im}\{F(z, z) + \mathbf{i}H(z, z)\}) dA(z)$
- (2). $-Q(A, A) \leq 16 \cdot (\int_X |F(z, z)|^2 dA(z) + \int_X |H(z, z)|^2 dA(z)).$

Proof. We begin with the third term in equation (3.4). By the Cauchy-Schwarz inequality, we may write

$$\begin{aligned}
(3.5) \quad & \left| \int_{X \times X} G(z, w)(F(z, w) + \mathbf{i}H(z, w))(F(w, z) + \mathbf{i}H(w, z)) dA(w) dA(z) \right| \\
& \leq \int_{X \times X} G(z, w) |(F(z, w) + \mathbf{i}H(z, w))(F(w, z) + \mathbf{i}H(w, z))| dA(w) dA(z) \\
& \leq \sqrt{\int_{X \times X} G(z, w) |(F(z, w) + \mathbf{i}H(z, w))|^2 dA(w) dA(z)} \\
& \quad \times \sqrt{\int_{X \times X} G(z, w) |(F(w, z) + \mathbf{i}H(w, z))|^2 dA(w) dA(z)} \\
& = \int_{X \times X} G(z, w) |(F(z, w) + \mathbf{i}H(z, w))|^2 dA(w) dA(z).
\end{aligned}$$

The last equality follows from the symmetry $G(z, w) = G(w, z)$.

It directly follows from inequality (3.5) that

$$\begin{aligned}
(3.6) \quad & \operatorname{Re} \int_{X \times X} G(z, w)(F(z, w) + \mathbf{i}H(z, w))(F(w, z) + \mathbf{i}H(w, z)) dA(w) dA(z) \\
& \leq \int_{X \times X} G(z, w) |(F(z, w) + \mathbf{i}H(z, w))|^2 dA(w) dA(z).
\end{aligned}$$

Recall that Q is negative semi-definite. Part (1) of the conclusion follows from the inequality above: the right hand side estimate above for the third term in equation (3.4) cancels with the second term in equation (3.4).

For part (2), beginning from the triangle inequality, we find

$$\begin{aligned}
(3.7) \quad |Q(A, A)| &\leq 4 \int_X D(\operatorname{Im}\{F(z, z) + \mathbf{i}H(z, z)\}) \cdot (\operatorname{Im}\{F(z, z) + \mathbf{i}H(z, z)\}) dA(z) \\
&\quad + 2 \int_{X \times X} G(z, w) |F(z, w) + \mathbf{i}H(z, w)|^2 dA(w) dA(z) \\
&\quad + 2 |\operatorname{Re} \int_{X \times X} G(z, w) (F(z, w) + \mathbf{i}H(z, w))(F(w, z) + \mathbf{i}H(w, z)) dA(w) dA(z)| \\
&\leq 4 \int_X D(\operatorname{Im}\{F(z, z) + \mathbf{i}H(z, z)\}) \cdot (\operatorname{Im}\{F(z, z) + \mathbf{i}H(z, z)\}) dA(z) \\
&\quad + 4 \int_{X \times X} G(z, w) |F(z, w) + \mathbf{i}H(z, w)|^2 dA(w) dA(z) \quad (\text{by applying (3.6)}) \\
&\leq 4 \int_X (\operatorname{Im}\{F(z, z) + \mathbf{i}H(z, z)\})^2 dA(z) \\
&\quad + 4 \int_{X \times X} G(z, w) |F(z, w) + \mathbf{i}H(z, w)|^2 dA(w) dA(z) \quad (\text{by Lemma 2.2}) \\
&\leq 4 \int_X |F(z, z) + \mathbf{i}H(z, z)|^2 dA(z) \\
&\quad + 4 \int_X D(|F(z, z) + \mathbf{i}H(z, z)|^2) dA(z)
\end{aligned}$$

where we apply part (1) of Proposition 2.3 for the last step.

Since D is self-adjoint and $D(1) = 1$, we see that we may rewrite the final term of (3.7),

$$\begin{aligned}
\int_X D(|F(z, z) + \mathbf{i}H(z, z)|^2) dA(z) &= \int_X |F(z, z) + \mathbf{i}H(z, z)|^2 \cdot D(1) dA(z) \\
&= \int_X |F(z, z) + \mathbf{i}H(z, z)|^2 dA(z).
\end{aligned}$$

This last equality allows us to combine the two terms on the right-hand side of the final inequality of (3.7) to obtain

$$\begin{aligned}
|Q(A, A)| &\leq 8 \int_X (|F(z, z) + \mathbf{i}H(z, z)|^2) dA(z) \\
&\leq 16 \cdot \left(\int_X |F(z, z)|^2 dA(z) + \int_X |H(z, z)|^2 dA(z) \right),
\end{aligned}$$

with the last inequality following from the elementary inequality

$$|z_1 + \mathbf{i}z_2|^2 \leq 2(|z_1|^2 + |z_2|^2), \quad \forall z_1, z_2 \in \mathbb{C}.$$

□

We use the first part of this proposition to prove Theorem 1.1 in the next section, and the second part of the proposition to prove Theorem 1.2 in Section 5.

4. A UNIFORM UPPER BOUND FOR $\lambda_{\min}(X)$

From the definition of the curvature operator we know that

$$\sum_{i=1}^{(3g-3)^2} \lambda_i(X) = Sca(X).$$

Thus a trivial upper bound for $\lambda_{(3g-3)^2}(X)$ is $\lambda_{(3g-3)^2}(X) \leq \frac{Sca(X)}{(3g-3)^2}$. By Remark 2.6 in section 2, this upper bound is less than or equal to $\frac{-1}{12\pi(g-1)}$, which approaches zero as g goes to infinity. The goal of this section is to prove Theorem 1.1 which estimates $\lambda_{(3g-3)^2}(X)$ by a negative number independent of genus.

The proof is actually quite straightforward: because $\lambda_{\min}(X)$ will measure the L^∞ -norm of \tilde{Q} , we simply exhibit an (unit normed) element A_0 in $\wedge^2 T_X \text{Teich}(S_g)$ with L^∞ norm bounded away from zero. After a few preparatory remarks on how one estimates the norm of \tilde{Q} , we display our element A_0 and verify the claims as to its \tilde{Q} -norm.

There are three terms in the formula of the Weil-Petersson curvature operator presented in Proposition 3.5. The first term is non-positive, and the sum of the second and third terms is also non-positive by the argument in the proof of Proposition 3.6. Of course, since $\lambda_{(3g-3)^2}(X)$ is the minimum eigenvalue, the Rayleigh-Ritz formulation implies that for any element A , we have that $\lambda_{(3g-3)^2}(X) \leq \frac{Q(A,A)}{\|A\|_{eu}^2}$. Now we are ready to describe the proof of Theorem 1.1 in detail.

Proof of Theorem 1.1. Let $\{\mu_i\}_{i=1}^{3g-3}$ be a holomorphic orthonormal basis $T_X \text{Teich}(S_g)$ and let $\{\frac{\partial}{\partial t_i}\}_{i=1}^{3g-3}$ be the vector field on $\text{Teich}(S_g)$ near X such that $\frac{\partial}{\partial t_i}|_X = \mu_i$. Let $t_i = x_i + \mathbf{i}y_i$ and let $A_0 = \frac{1}{\sqrt{3g-3}}(\sum_{i=1}^{3g-3} \frac{\partial}{\partial x_i}(X) \wedge \frac{\partial}{\partial y_i}(X))$ be our special element of $\wedge^2 T_X \text{Teich}(S_g)$.

We begin with some preliminary computations on our special two-form A_0 : we show that $\|A_0\|_{eu} = 1$, and find the associated functions F and H used in the formula (3.4) for the associated form $Q(A_0, A_0)$ for this element. Since $\{\mu_i\}_{i=1}^{3g-3}$ is a holomorphic orthonormal basis $T_X \text{Teich}(S_g)$, we have

$$\begin{aligned} \langle \frac{\partial}{\partial x_i}(X), \frac{\partial}{\partial x_j}(X) \rangle &= \text{Re}\{\langle \frac{\partial}{\partial x_i}(X) + \mathbf{i}\mathbf{J} \circ \frac{\partial}{\partial x_i}(X), \frac{\partial}{\partial x_j}(X) + \mathbf{i}\mathbf{J} \circ \frac{\partial}{\partial x_j}(X) \rangle\} \\ &= \text{Re}\{\langle \mu_i, \mu_j \rangle\} = \delta_{ij}. \end{aligned}$$

Similarly, we have

$$\begin{aligned} \langle \frac{\partial}{\partial y_i}(X), \frac{\partial}{\partial y_j}(X) \rangle &= \text{Re}\{\langle \frac{\partial}{\partial y_i}(X) + \mathbf{i}\mathbf{J} \circ \frac{\partial}{\partial y_i}(X), \frac{\partial}{\partial y_j}(X) + \mathbf{i}\mathbf{J} \circ \frac{\partial}{\partial y_j}(X) \rangle\} \\ &= \text{Re}\{\langle -\mathbf{i}\mu_i, -\mathbf{i}\mu_j \rangle\} = \text{Re}\{\langle \mu_i, \mu_j \rangle\} = \delta_{ij} \end{aligned}$$

and

$$\begin{aligned}
\left\langle \frac{\partial}{\partial x_i}(X), \frac{\partial}{\partial y_j}(X) \right\rangle &= \operatorname{Re}\left\{ \left\langle \frac{\partial}{\partial x_i}(X) + \mathbf{i}\mathbf{J} \circ \frac{\partial}{\partial x_i}(X), \frac{\partial}{\partial x_i}(X) + \mathbf{i}\mathbf{J} \circ \frac{\partial}{\partial y_j}(X) \right\rangle \right\} \\
&= \operatorname{Re}\left\{ \langle \mu_i, -\mathbf{i}\mu_j \rangle \right\} = -\operatorname{Im}\left\{ \langle \mu_i, \mu_j \rangle \right\} \\
&= -\operatorname{Im}\{\delta_{ij}\} = 0.
\end{aligned}$$

Thus, equation (3.1) gives that

$$\begin{aligned}
&\left\langle \frac{\partial}{\partial x_i}(X) \wedge \frac{\partial}{\partial y_i}(X), \frac{\partial}{\partial x_j}(X) \wedge \frac{\partial}{\partial y_j}(X) \right\rangle_{eu} \\
&= \left\langle \frac{\partial}{\partial x_i}(X), \frac{\partial}{\partial x_j}(X) \right\rangle \left\langle \frac{\partial}{\partial y_i}(X), \frac{\partial}{\partial y_j}(X) \right\rangle \\
&- \left\langle \frac{\partial}{\partial x_i}(X), \frac{\partial}{\partial y_j}(X) \right\rangle \left\langle \frac{\partial}{\partial y_i}(X), \frac{\partial}{\partial x_j}(X) \right\rangle \\
&= \delta_{ij}^2 - 0 = \delta_{ij}.
\end{aligned}$$

Therefore, the norm of A_0 satisfies

$$(4.1) \quad \|A_0\|_{eu}^2 = \frac{1}{3g-3} \left\langle \sum_{i=1}^{3g-3} \frac{\partial}{\partial x_i}(X) \wedge \frac{\partial}{\partial y_i}(X), \sum_{i=1}^{3g-3} \frac{\partial}{\partial x_i}(X) \wedge \frac{\partial}{\partial y_i}(X) \right\rangle_{eu} = 1.$$

As we plan to apply Proposition 3.6, we need to compute $F(z, z)$ and $H(z, z)$ for this element A_0 . Of course,

$$F(z, z) = \sum_{i,j=1}^{3g-3} (a_{ij} + c_{ij}) \mu_i(z) \cdot \overline{\mu_j(z)} = 0$$

as from our definition of A_0 and the coefficients a_{ij}, c_{ij} in (3.3), we see we have set all of the $a_{ij} = c_{ij} = 0$ in the definition of $F(z, z)$ in Proposition 3.5. Similarly,

$$H(z, w) = \sum_{i,j=1}^{3g-3} b_{ij} \mu_i(w) \cdot \overline{\mu_j(z)} = \sum_{i=1}^{3g-3} \frac{1}{\sqrt{3g-3}} \mu_i(w) \cdot \overline{\mu_i(z)}$$

by our definition of H in Proposition 3.5.

From part (1) of Proposition 3.6,

$$\begin{aligned}
Q(A_0, A_0) &\leq -4 \int_X D(\operatorname{Im}\{\mathbf{i}H(z, z)\}) \cdot (\operatorname{Im}\{\mathbf{i}H(z, z)\}) dA(z) \\
&= -4 \int_X D\left(\frac{1}{\sqrt{3g-3}} \sum_{i=1}^{3g-3} |\mu_i|^2\right) \left(\frac{1}{\sqrt{3g-3}} \sum_{i=1}^{3g-3} |\mu_i|^2\right) dA \\
&\leq -\frac{4}{3} \int_X \left(\frac{1}{\sqrt{3g-3}} \sum_{i=1}^{3g-3} |\mu_i|^2\right)^2 dA \quad (\text{by Lemma 2.1}) \\
&= -\frac{4}{9(g-1)} \int_X \left(\sum_{i=1}^{3g-3} |\mu_i|^2\right)^2 dA \\
&\leq \frac{2}{9(g-1)} \operatorname{Sca}(X),
\end{aligned}$$

with the last inequality follows from Proposition 2.5.

Combining equation (3.2), equation (4.1) and the inequality above, we have

$$\lambda_{\min}(X) \leq \frac{Q(A_0, A_0)}{\|A_0\|_{eu}^2} \leq \frac{2\operatorname{Sca}(X)}{9(g-1)}.$$

By Lemma 4.6 in [34], Wolpert's upper bound for scalar curvature, we have

$$\lambda_{\min}(X) \leq \frac{2\operatorname{Sca}(X)}{9(g-1)} \leq \frac{2}{9(g-1)} \cdot \frac{-3(3g-2)}{4\pi} < \frac{-1}{2\pi}.$$

□

5. A UNIFORM LOWER BOUND FOR $\lambda_{\min}(X)$ IN THE THICK-PART

The goal of this section is to prove Theorem 1.2 that was stated in the introduction. In contrast to the argument in the last section where we bounded from above the norm of a particular element $A_0 \in \wedge^2 T_X \operatorname{Teich}(S_g)$, here we need to bound from below the norm of an arbitrary element $A \in \wedge^2 T_X \operatorname{Teich}(S_g)$. The proof rests on the (other) bound in Proposition 3.5, where here we inherit uniform bounds on all of the terms in the the norm $Q(A, A)$ from the uniform thickness of a surface in the thick part of the surface.

Recall the systole, denoted by $\operatorname{systole}(X)$, of a compact hyperbolic surface X is the length of the shortest nontrivial simple closed curve in X . So $\operatorname{systole}(X) = 2 \operatorname{inj}(X)$ where $\operatorname{inj}(X)$ is the injectivity radius of X . The ϵ -thick part $\operatorname{Teich}(S_g)^{\geq \epsilon}$ of $\operatorname{Teich}(S_g)$ is defined by

$$\operatorname{Teich}(S_g)^{\geq \epsilon} := \{Y \in \operatorname{Teich}(S_g); \operatorname{systole}(Y) \geq \epsilon\}.$$

Let $\operatorname{Mod}(S)$ be the mapping class group of S . Then the quotient space $\mathbb{M}(S)^{\geq \epsilon} := \operatorname{Teich}(S_g)^{\geq \epsilon} / \operatorname{Mod}(S)$ is called the ϵ -thick part of the moduli space. By Mumford compactness, we have (see e.g. [12]) that $\mathbb{M}(S)^{\geq \epsilon}$ is compact for all $\epsilon > 0$. In particular there is a lower bound for $\lambda_{(3g-3)^2}(X)$

where X runs over the thick part of $\text{Teich}(S_g)$. A natural question is whether this lower bound depends on the topology of the surface. Theorem 1.2 answers this question negatively; specifically, we provide a lower bound that only depends on the thickness ϵ but not on genus of the surface.

We begin the proof of Theorem 1.2 with some preliminary estimates on norms of sums of harmonic Beltrami differentials. Let $\{\mu_i\}_{i=1}^{3g-3}$ be a holomorphic orthonormal basis of $T_X \text{Teich}(S_g)$; thus, in particular, we have that $\int_X \mu_i(z) \overline{\mu_j(z)} dA = \delta_{ij}$. Now, the expression of the Weil-Petersson curvature operator – presented in Proposition 3.5 – involved expressions $F(z, w)$ and $H(z, w)$ both of the same form, say $K(z, w) = \sum_{i,j=1}^{3g-3} d_{ij} \mu_i(w) \cdot \overline{\mu_j(z)}$ (where the coefficients d_{ij} were real). We separate the proof of Theorem 1.2 into lemmas that estimate norms of such expressions $K(z, w)$. The following lemma is a direct consequence of Proposition 2.10.

Lemma 5.1. *Fix $z \in X$, we have*

$$\sup_{w \in X} |K(z, w)|^2 \leq C(\text{inj}(X)) \cdot \left(\sum_{1 \leq i, j, l \leq 3g-3} d_{ij} d_{il} \overline{\mu_j(z)} \mu_l(z) \right)$$

where $C(\text{inj}(X))$ is the same as in Proposition 2.10.

Proof. Rewrite $K(z, w)$ as

$$K(z, w) = \sum_{i=1}^{3g-3} \left(\sum_{j=1}^{3g-3} d_{ij} \overline{\mu_j(z)} \right) \cdot \mu_i(w).$$

Thus, if we fix $z \in X$, the form $K(z, w)$ is a harmonic Beltrami differential on X in the coordinate w . From Proposition 2.10, we have

$$\begin{aligned} \sup_{w \in X} |K(z, w)|^2 &\leq C(\text{inj}(X)) \cdot \int_X K(z, w) \overline{K(z, w)} dA(w) \\ &= C(\text{inj}(X)) \cdot \int_X \left(\sum_{i,j=1}^{3g-3} d_{ij} \mu_i(w) \cdot \overline{\mu_j(z)} \right) \left(\sum_{k,l=1}^{3g-3} d_{kl} \overline{\mu_k(w)} \cdot \mu_l(z) \right) dA(w) \\ &= C(\text{inj}(X)) \cdot \left(\sum_{1 \leq i, j, k, l \leq 3g-3} d_{ij} d_{kl} \overline{\mu_j(z)} \mu_l(z) \right) \cdot \int_X \mu_i(w) \overline{\mu_k(w)} dA(w) \\ &= C(\text{inj}(X)) \cdot \left(\sum_{1 \leq i, j, l \leq 3g-3} d_{ij} d_{il} \overline{\mu_j(z)} \mu_l(z) \right), \end{aligned}$$

where the last equality uses the fact that the basis $\{\mu_i\}$ is orthonormal. \square

Specializing to an L^2 bound for $K(z, z)$, we find

Lemma 5.2. $\int_X |K(z, z)|^2 dA(z) \leq C(\text{inj}(X)) \cdot \left(\sum_{1 \leq i, j \leq (3g-3)} d_{ij}^2 \right)$.

Proof. First, Lemma 5.1 gives that, for any $z \in X$,

$$|K(z, z)|^2 \leq \sup_{w \in X} |K(z, w)|^2 \leq C(\text{inj}(X)) \cdot \left(\sum_{1 \leq i, j, l \leq 3g-3} d_{ij} d_{il} \overline{\mu_j(z)} \mu_l(z) \right).$$

Thus, we have

$$\begin{aligned}
& \int_X |K(z, z)|^2 dA(z) \\
& \leq C(\text{inj}(X)) \cdot \int_X \sum_{1 \leq i, j, l \leq 3g-3} d_{ij} d_{il} \overline{\mu_j(z)} \mu_l(z) dA(z) \\
& = C(\text{inj}(X)) \cdot \left(\sum_{1 \leq i, j \leq (3g-3)} d_{ij}^2 \right)
\end{aligned}$$

where the last equality follows from the assumption that $\{\mu_i\}$ is orthonormal. \square

Now we are ready to prove Theorem 1.2.

Proof of Theorem 1.2. Let $A \in \wedge^2 T_X \text{Teich}(S_g)$ be expressed as

$$A = \left(\sum_{1 \leq i < j \leq (3g-3)} a_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j} + \sum_{1 \leq i, j \leq (3g-3)} b_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j} + \sum_{1 \leq i < j \leq (3g-3)} c_{ij} \frac{\partial}{\partial y_i} \wedge \frac{\partial}{\partial y_j} \right)$$

where a_{ij}, b_{ij}, c_{ij} are real. In terms of the coefficients of the basic forms, we may compute the norm $\|A\|_{eu}$ of A as

$$(5.1) \quad \|A\|_{eu}^2 = \sum_{1 \leq i < j \leq (3g-3)} a_{ij}^2 + \sum_{1 \leq i, j \leq (3g-3)} b_{ij}^2 + \sum_{1 \leq i < j \leq (3g-3)} c_{ij}^2.$$

From part (2) of Proposition 3.6, we have

$$Q(A, A) \geq -16 \cdot \left(\int_X |F(z, z)|^2 dA(z) + \int_X |H(z, z)|^2 dA(z) \right)$$

where $F(z, z) = \sum_{1 \leq i < j \leq (3g-3)} (a_{ij} + c_{ij}) \mu_i(z) \cdot \overline{\mu_j(z)}$ and $H(z, z) = \sum_{i, j=1}^{3g-3} b_{ij} \mu_i(z) \cdot \overline{\mu_j(z)}$.

By Lemma 5.2, we have

$$\int_X |F(z, z)|^2 dA(z) \leq C(\text{inj}(X)) \cdot \left(\sum_{1 \leq i < j \leq (3g-3)} (a_{ij} + c_{ij})^2 \right)$$

where we have set $\{d_{ij} = a_{ij} + c_{ij}\}$ for $i < j$ and $d_{ij} = 0$ otherwise.

Similarly, by Lemma 5.2, we also have

$$\int_X |H(z, z)|^2 dA(z) \leq C(\text{inj}(X)) \cdot \left(\sum_{1 \leq i, j \leq (3g-3)} (b_{ij})^2 \right)$$

by setting $\{d_{ij} = b_{ij}\}$.

Combine these two inequalities above, we find

$$\begin{aligned}
Q(A, A) & \geq -16C(\text{inj}(X)) \cdot \left(\sum_{1 \leq i < j \leq (3g-3)} (a_{ij} + c_{ij})^2 + \sum_{1 \leq i, j \leq (3g-3)} b_{ij}^2 \right) \\
& \geq -32C(\text{inj}(X)) \cdot \left(\sum_{1 \leq i < j \leq (3g-3)} (a_{ij}^2 + c_{ij}^2) + \sum_{1 \leq i, j \leq (3g-3)} b_{ij}^2 \right) \\
& = -32C(\text{inj}(X)) \|A\|_{eu}^2 \text{ by (5.1)}.
\end{aligned}$$

The theorem then follows from the Rayleigh-Ritz characterization of the lowest eigenvalue and by choosing $B(\epsilon) = 32C(\frac{\epsilon}{2})$. \square

Remark 5.3. Theorem 1.2 shows the Weil-Petersson curvature operators, restricted to the thick part of the moduli space, are uniformly bounded by a constant that depends only on the bound on prescribed thickness (the lower bound in injectivity radius) of the surfaces in the given thick part, but not on the topology of the underlying surface. It is worth noting that the constant $B(\epsilon) \asymp \frac{1}{\epsilon^2}$ tends to infinity as $\epsilon \rightarrow 0$. Of course, it is known that sectional curvatures (the diagonal elements for the curvature operator \tilde{Q}) decay at worst with order $O(\frac{-1}{\epsilon})$. (One can see [9] and [39] for details.) We do not know if this bound of $O(\frac{-1}{\epsilon})$, sharp in the case of sectional curvatures, also extends to the broader setting of the curvature operator. On the other hand, as $\epsilon \rightarrow \infty$, the constant $B(\epsilon) \rightarrow \frac{24}{\pi} > \frac{1}{2\pi}$, which is consistent with the bound we found for a special element of $\wedge^2 T_X \text{Teich}(S_g)$ in Theorem 1.1.

Let R be the Riemannian curvature tensor of the Weil-Petersson metric. Let $X \in \text{Teich}(S_g)$ and $\{e_i\}_{i=1}^{6g-6}$ be an orthonormal basis of $T_X \text{Teich}(S_g)$. Recall that $R_{ijkl} := \langle R(e_i, e_j)e_k, e_l \rangle$. Define the norm $\|R\|_X$ of the curvature tensor R on X by

$$\|R\|_X = \sup_{1 \leq i, j, k, l \leq (6g-6)} \max |R_{ijkl}|$$

where the supremum runs over all the orthonormal bases of $T_X \text{Teich}(S_g)$.

A simple application of Theorem 1.2 is

Corollary 5.4. *Let $X \in \text{Teich}(S_g)$ and R be the Riemannian curvature tensor of the Weil-Petersson metric. Then,*

$$\|R\|_X \leq \hat{B}(\text{inj}(X))$$

where $\text{inj}(X)$ is the injectivity radius of X and $\hat{B}(\text{inj}(X))$ is a function of $\text{inj}(X)$.

Proof. Set $\|Q(X)\| := \sup_{A \in \wedge^2 T_X \text{Teich}(S_g), \|A\|_{eu}=1} |\langle \tilde{Q}(A), A \rangle|$. Theorem 1.2 gives that $\|Q(X)\| \leq B(2 \text{inj}(X))$. Let $\{e_i\}_{i=1}^{6g-6}$ be an orthonormal basis of $T_X \text{Teich}(S_g)$. Then the Cauchy Schwarz inequality leads to

$$\begin{aligned} |R_{ijkl}| &= |\langle \tilde{Q}(e_i \wedge e_j), e_k \wedge e_l \rangle| \\ &\leq |\tilde{Q}(e_i \wedge e_j)| \cdot |e_k \wedge e_l| \leq \|Q(X)\| \leq B(2 \text{inj}(X)). \end{aligned}$$

Since the indices $\{i, j, k, l\}$ are arbitrary, the conclusion follows by choosing $\hat{B}(\text{inj}(X)) = B(2 \text{inj}(X))$. \square

Remark 5.5. For the case that $\{i = k, j = l\}$, R_{ijij} is the sectional curvature (using the orthonormality of the coordinate system). Huang in [10] proved the the sectional curvature, restricted on the thick part of the moduli space, is uniformly bounded from below by using harmonic maps.

The Weil-Petersson metric is not complete (see [7, 33]). We let $\overline{\text{Teich}(S_g)}$ be the metric completion of $\text{Teich}(S_g)$ and $\partial\overline{\text{Teich}(S_g)}$ be the frontier of $\overline{\text{Teich}(S_g)}$: this frontier is composed of products of lower dimensional Teichmüller spaces (one can see [19, 36, 38] for more details). For any $X \in \text{Teich}(S_g)$, the following quantitative version of estimation on the distance between X and $\partial\overline{\text{Teich}(S_g)}$ is provided by Wolpert.

Theorem 5.6 ([37]). $\text{dist}(X, \partial\overline{\text{Teich}(S_g)}) \leq \sqrt{4\pi \cdot \text{inj}(X)}$.

The following consequence is motivated by Proposition 4.22 in [3].

Corollary 5.7. *There exists a constant $C > 0$, which is independent of the topology of the surface, such that the norm $\|R\|_X$ of the curvature tensor R on X satisfies*

$$\|R\|_X \leq \max\left\{C, \frac{C}{\text{dist}(X, \partial\overline{\text{Teich}(S_g)})^4}\right\}.$$

Proof. We observed after stating Proposition 2.10 that as $\epsilon \rightarrow 0$, the constant $C(\epsilon)$ is asymptotic to $\frac{1}{\pi\epsilon^2}$. Fix $\epsilon_0 > 0$ such that $C(\epsilon) \leq \frac{2}{\pi\epsilon^2}$ for all $\epsilon \in (0, \epsilon_0]$. Then for any $X \in \text{Teich}(S_g)$, there are two possibilities:

Case 1: If the injectivity radius $\text{inj}(X) \geq \epsilon_0$, the inequality $\|R\|_X \leq \widehat{B}(\epsilon_0) = B(2\epsilon_0)$ follows from Corollary 5.4.

Case 2: If the injectivity radius $\text{inj}(X) \leq \epsilon_0$, then from Corollary 5.4 we find,

$$\|R\|_X \leq 32C(\text{inj}(X)) \leq \frac{64}{\pi \cdot \text{inj}(X)^2} \leq \frac{1024\pi}{\text{dist}(X, \partial\overline{\text{Teich}(S_g)})^4},$$

where we apply Theorem 5.6 for the last inequality. The conclusion follows by choosing $C = \max\{B(2\epsilon_0), 1024\pi\}$. \square

Remark 5.8. We emphasize here that the constant C is independent not only of the genus g , but also of any choice of neighborhood of $\partial\overline{\text{Teich}(S_g)}$.

6. EIGENVALUES OF THE WEIL-PETERSSON CURVATURE OPERATORS ON THICK SURFACES

Our goal in this final section of this part of the paper is to prove the remaining results, Theorem 1.3 and Theorem 1.5, on the Weil-Petersson curvature operator. The proofs are reasonably straightforward consequences of the results and tools we have already developed.

We first prove Theorem 1.3 which bounds the smallest eigenvalue λ_{\min} of the Weil-Petersson curvature operator on surfaces of sufficiently large genus whose injectivity radii are increasing without bound.

Proof of Theorem 1.3. First, one computes that the constant $B(\epsilon)$ in Theorem 1.2 satisfies

$$\lim_{\epsilon \rightarrow \infty} B(\epsilon) = \frac{24}{\pi}.$$

The conclusion then follows from Theorem 1.1, Theorem 1.2 and the limit above. \square

Next, fix $\epsilon > 0$; then the Mumford compactness theorem implies the ϵ -thick part $\mathbb{M}^{\geq \epsilon}$ of the moduli space of a closed surface is compact. Thus the following functions are well-defined:

$$\underline{\lambda}_i^\epsilon(g) := \min_{X \in \text{Teich}(S_g)^{\geq \epsilon}} \lambda_i(X), \quad \forall 1 \leq i \leq (3g-3)^2$$

and

$$\overline{\lambda}_i^\epsilon(g) := \max_{X \in \text{Teich}(S_g)^{\geq \epsilon}} \lambda_i(X), \quad \forall 1 \leq i \leq (3g-3)^2.$$

Of course

$$\underline{\lambda}_i^\epsilon(g) \leq \overline{\lambda}_i^\epsilon(g),$$

and Theorems 1.1 and 1.2 imply

$$-B(\epsilon) \leq \underline{\lambda}_{(3g-3)^2}^\epsilon(g) \leq \overline{\lambda}_{(3g-3)^2}^\epsilon(g) \leq \frac{-1}{2\pi}.$$

Our focus for the rest of this section then turns from the smallest eigenvalue $\lambda_{(3g-3)^2} = \lambda_{\min}$ to the asymptotic properties for $\underline{\lambda}_i^\epsilon(g)$ and $\overline{\lambda}_i^\epsilon(g)$ as g goes to infinity where the index i is not maximal but instead small.

Let

$$f : \{1, 2, \dots, (3g-3)^2\} \rightarrow \{1, 2, \dots, (3g-3)^2\}$$

be a function. Recall that Theorem 1.5 states that if the index i is not close to the maximal value $(3g-3)^2$ in the limit sense, then the bound $\underline{\lambda}_i^\epsilon(g)$ tends to zero as g goes to infinity.

Proof of Theorem 1.5. Since $1 \leq f(g) \leq (3g-3)^2$, It follows from Theorem 3.3 that $\overline{\lambda}_{f(g)}^\epsilon(g) < 0$. For the lower bound, let $X \in \text{Teich}(S_g)^{\geq \epsilon}$ and recall that $\{\lambda_i(X)\}_{i=1}^{(3g-3)^2}$ is the set of the non-zero eigenvalues of the Weil-Petersson curvature operator at X with $\lambda_{i+1}(X) \leq \lambda_i(X)$. Proposition 3.3 in [29] tells us that

$$-6(g-1)C\left(\frac{\epsilon}{2}\right) \leq \text{Sca}(X) = \sum_{i=1}^{(3g-3)^2} \lambda_i(X),$$

where $C(\frac{\epsilon}{2})$ is defined in Proposition 2.10. Since $\lambda_i(X) < 0$, we have that

$$-6(g-1)C\left(\frac{\epsilon}{2}\right) \leq \sum_{i=f(g)}^{(3g-3)^2} \lambda_i(X) \leq ((3g-3)^2 - f(g) + 1)\lambda_{f(g)}(X).$$

We rewrite this last inequality as

$$\frac{-6C(\frac{\epsilon}{2})}{g} \cdot \frac{g(g-1)}{(3g-3)^2 - f(g) + 1} \leq \lambda_{f(g)}(X).$$

Then, since $\limsup_{g \rightarrow \infty} \frac{g(g-1)}{(3g-3)^2 - f(g) + 1} = \frac{1}{9(1-\alpha)} > 0$ and $X \in \text{Teich}(S_g)^{\geq \epsilon}$ is arbitrary, the conclusion follows by choosing $B(\alpha, \epsilon) = \frac{4C(\frac{\epsilon}{2})}{3(1-\alpha)}$. \square

Remark 6.1. (1). Recall that as $\epsilon \rightarrow 0$, the constant $C(\epsilon)$ behaves like

$$\lim_{\epsilon \rightarrow 0} \frac{C(\epsilon)}{\frac{1}{\pi\epsilon^2}} = 1.$$

The proof of Theorem 1.5 then yields the following more general statement.

Theorem 6.2. *If the function f satisfies $\limsup_{g \rightarrow \infty} \frac{f(g)}{9g^2} = \alpha < 1$. Let X_g be a sequence of Riemannian surfaces whose injectivity radii satisfy $\lim_{g \rightarrow \infty} \frac{1}{g \cdot \text{inj}(X_g)^2} = 0$. Then,*

$$\lim_{g \rightarrow \infty} \lambda_{f(g)}(X_g) = 0.$$

This theorem tells us that as the genus goes to infinity, even in certain portions of the thin parts of the moduli spaces, we could still have

$$\lim_{g \rightarrow \infty} \bar{\lambda}_i^{\epsilon(g)}(g) = \lim_{g \rightarrow \infty} \lambda_i^{\epsilon(g)}(g) = 0, \quad \text{for all } 1 \leq i \leq 8g^2$$

provided that $\lim_{g \rightarrow \infty} \frac{1}{g \cdot \epsilon(g)^2} = 0$.

(2). *We do not know any asymptotic properties of $\lambda_i^\epsilon(g)$ and $\bar{\lambda}_i^\epsilon(g)$ as g goes to infinity when i is close to $(3g-3)^2$; for example $\lambda_{(3g-3)^2-g}^\epsilon(g)$ and $\bar{\lambda}_{(3g-3)^2-g}^\epsilon(g)$. It is interesting to study them.*

Part 2. Bounds on the Weil-Petersson holomorphic sectional curvatures

7. UNIFORM ESTIMATES ON HARMONIC BELTRAMI DIFFERENTIALS.

Recall from section 2.3 that a surface is C -sufficiently thick if $\text{inj}(X) \geq C$, and that there are non-trivial subspaces of the Teichmüller spaces $\text{Teich}(S_g)$ composed of sufficiently thick surfaces, once the genus g is chosen sufficiently large. In this section will construct holomorphic lines in the tangent spaces of sufficiently thick surfaces (of course, necessarily of large genera and injectivity radii) such that those sections have uniformly pinched negative holomorphic sectional curvatures. The pinching constants will be independent of the genus.

To begin, we pick up the thread on which we ended section 2.2.2, that related holomorphic sectional curvatures to powers of normalized L^∞ norms of harmonic Beltrami differentials. In particular, we will directly apply Proposition 2.11 to a specific choice of harmonic Beltrami differential μ_g , whose construction we begin immediately below. As we noted at the end of section 2.2.2, our (main) goal is to bound the L^2 norm of μ_g from above and the L^∞ norm of μ_g from below.

7.1. Poincaré series. Let $X_g \in \text{Teich}(S_g)$ and let $\Gamma_g \subset \text{Aut}(\mathcal{D})$ be a representation of $\pi_1(X_g)$ into the (Möbius) group $\text{Aut}(\mathcal{D})$ of automorphisms of the hyperbolic disk \mathcal{D} where $\pi : \mathcal{D} \rightarrow X_g$ is the universal covering map. Let $H(\mathcal{D})$ denote the set of holomorphic functions on the unit disk \mathcal{D} and $H^0(X_g, K^2)$ be the space of holomorphic quadratic differentials on X_g .

Recall the Theta-operator Θ is defined as

$$\begin{aligned} \Theta : H(\mathcal{D}) &\rightarrow H^0(X_g, K^2) \\ f(z) &\mapsto \sum_{\gamma \in \Gamma_g} f(\gamma(z)) \cdot \gamma'(z)^2. \end{aligned}$$

It is well-known that $\Theta(f)(z)$ is well-defined if f is integrable on \mathcal{D} , and this operator is surjective on its domain. (See Theorem 7.2 in [12] for details.)

Fix the constant function $f(z) \equiv 1$ be on \mathcal{D} . Then

$$\Theta(f(z)) = \Theta(1)(z) = \sum_{\gamma \in \Gamma_g} \gamma'(z)^2$$

whose corresponding harmonic Beltrami differential is

$$(7.1) \quad \mu_g(z) := \frac{\overline{\Theta(1)(z)}}{\rho(z)} = \sum_{\gamma \in \Gamma_g} \frac{\overline{\gamma'(z)^2}}{\rho(z)}$$

where $\rho(z) = \frac{4}{(1-|z|^2)^2}$ is the hyperbolic metric on the disk.

Ahlfors in [2] showed that $|\mu_g|$ is bounded above by a constant depending on Γ_g . In this section we will observe that $|\mu_g|$ is uniformly bounded above if we assume that the injectivity radius $\text{inj}(X_g) \rightarrow \infty$ as $g \rightarrow \infty$.

Let us close this subsection by recalling Ahlfors' method in [2], a technique which we will adapt for the heart of our argument.

Ahlfors' Method: From the triangle inequality we know that

$$(7.2) \quad |\mu_g(z)| \leq \sum_{\gamma \in \Gamma_g} \frac{|\gamma'(z)|^2}{\rho(z)}.$$

Then since $\rho(\gamma(z))|\gamma'(z)|^2 = \rho(z)$ for any $\gamma \in \Gamma_g$, and $\rho(\zeta) = 4(1-|\zeta|^2)^{-2}$, we have

$$(7.3) \quad \sum_{\gamma \in \Gamma_g} \frac{|\gamma'(z)|^2}{\rho(z)} = \frac{1}{4} \sum_{\gamma \in \Gamma_g} (1-|\gamma(z)|^2)^2.$$

Combining the above inequalities yields

$$(7.4) \quad |\mu_g(z)| \leq \frac{1}{4} \sum_{\gamma \in \Gamma_g} (1-|\gamma(z)|^2)^2.$$

Let Δ be the (Euclidean) Laplace operator on the (Euclidean) disk. Then a direct computation shows

$$\Delta\left(\sum_{\gamma \in \Gamma_g} (1 - |\gamma(z)|^2)^2\right) = 8 \cdot \sum_{\gamma \in \Gamma_g} (2|\gamma(z)|^2 - 1)|\gamma'(z)|^2.$$

Note that the terms on the right side are non-negative when $|\gamma(z)|^2 \geq \frac{1}{2}$. With that in mind, let $B(0, \frac{1}{\sqrt{2}}) := \{z \in \mathcal{D}; |z| \leq \frac{1}{\sqrt{2}}\}$ be the ball of Euclidean radius $\frac{1}{\sqrt{2}}$ and let $V := \cup_{\gamma \in \Gamma_g} \gamma^{-1} \circ B(0, \frac{1}{\sqrt{2}})$ be the pullbacks of this ball B by the group Γ_g . The equation above gives that $\sum_{\gamma \in \Gamma_g} (1 - |\gamma(z)|^2)^2$ is subharmonic in $\mathcal{D} - V$. Since both $\sum_{\gamma \in \Gamma_g} (1 - |\gamma(z)|^2)^2$ and V are Γ_g -invariant, and Γ_g is cocompact, we find

$$(7.5) \quad \sup_{z \in \mathcal{D}} \sum_{\gamma \in \Gamma_g} (1 - |\gamma(z)|^2)^2 = \sup_{z \in V} \sum_{\gamma \in \Gamma_g} (1 - |\gamma(z)|^2)^2 = \sup_{z \in B(0, \frac{1}{\sqrt{2}})} \sum_{\gamma \in \Gamma_g} (1 - |\gamma(z)|^2)^2$$

which in particular is bounded above by a constant depending on Γ_g : we display a version of the argument that the last term on the right is bounded in the next subsection: see especially equations (7.7) - (7.9).

7.2. Uniform upper bound for μ_g . Recall the relation between the Euclidean distance and the hyperbolic distance is

$$\text{dist}_{\mathcal{D}}(0, z) = \ln \frac{1 + |z|}{1 - |z|}.$$

For the rest of this section, we consider a surface X_g which is sufficiently thick – we retain the index g as a reminder that such surfaces have large genus g and we are computing curvatures of the Teichmüller space $\text{Teich}(S_g)$ with all of the normalizations that accompany that choice. (This will also be useful for a remark at the end of this section.) Therefore we may assume that we may find a ball $B(0, \frac{1}{\sqrt{2}} + \frac{1}{10}) \subset F_g$ where $F_g \subset \mathcal{D}$ is the (Dirichlet) fundamental domain, centered at the origin, of X_g . Applying the triangle inequality to equation (7.5), we have

$$(7.6) \quad \sup_{z \in \mathcal{D}} \sum_{\gamma \in \Gamma_g} (1 - |\gamma(z)|^2)^2 \leq 1 + \sup_{z \in B(0, \frac{1}{\sqrt{2}})} \sum_{\gamma \neq e} (1 - |\gamma(z)|^2)^2.$$

As in Ahlfors' method, by applying the Laplace operator, we obtain

$$(7.7) \quad \Delta\left(\sum_{\gamma \neq e} (1 - |\gamma(z)|^2)^2\right) = 8 \cdot \sum_{\gamma \neq e} (2|\gamma(z)|^2 - 1)|\gamma'(z)|^2.$$

Because of our assumption on the large injectivity radius of X_g , we see that $B(0, \frac{1}{\sqrt{2}} + \frac{1}{10}) \subset F_g$. Thus for $z \in B(0, \frac{1}{\sqrt{2}})$, we see that $\gamma(z) \notin B(0, \frac{1}{\sqrt{2}})$ when $\gamma \neq e$: in particular, for such γ , we have $|\gamma(z)|^2 \geq \frac{1}{2}$. This implies that for $z \in B(0, \frac{1}{\sqrt{2}} + \frac{1}{10})$ we have

$$(7.8) \quad \Delta \left(\sum_{\gamma \neq e} (1 - |\gamma(z)|^2)^2 \right) \geq 0.$$

Let $\zeta = \xi + i\eta$. Thus, by the mean-value inequality, for any $z \in B(0, \frac{1}{\sqrt{2}})$, we have

$$\begin{aligned} \sum_{\gamma \neq e} (1 - |\gamma(z)|^2)^2 &\leq \frac{1}{B(z, \frac{1}{10})} \int_{B(z, \frac{1}{10})} \sum_{\gamma \neq e} (1 - |\gamma(\zeta)|^2)^2 d\xi d\eta \\ &= \frac{100}{\pi} \int_{B(z, \frac{1}{10})} \sum_{\gamma \neq e} \frac{|\gamma'(\zeta)|^2}{\rho(\zeta)} d\xi d\eta \text{ by equation (7.3)} \\ &\leq \frac{25}{\pi} \int_{B(z, \frac{1}{10})} \sum_{\gamma \neq e} |\gamma'(\zeta)|^2 d\xi d\eta \text{ since } \rho(z) = 4(1 - |z|^2)^{-2} \geq 4 \\ &= \frac{25}{\pi} \int_{\cup_{\gamma \neq e} \gamma \circ B(z, \frac{1}{10})} d\xi d\eta \text{ after unfolding the sum} \\ &\leq \frac{25}{\pi} \int_{\mathcal{D} - F_g} d\xi d\eta \\ &= \frac{25}{\pi} (\pi - \text{Area}(F_g)). \end{aligned}$$

Since $z \in B(0, \frac{1}{\sqrt{2}})$ is arbitrary,

$$(7.9) \quad \sup_{z \in B(0, \frac{1}{\sqrt{2}})} \sum_{\gamma \neq e, \gamma \in \Gamma_g} (1 - |\gamma(z)|^2)^2 \leq \frac{25}{\pi} (\pi - \text{Area}(F_g)).$$

Finally, we invoke the hypothesis that the surface X_g is sufficiently thick. Assuming that thickness, we may assume that the Euclidean area $\text{Area}(F_g)$ of the fundamental domain F_g is as close to π as we wish, so the quantitative version of sufficiently thick that we will invoke will force

$$(7.10) \quad \frac{25}{\pi} (\pi - \text{Area}(F_g)) < \frac{1}{4}.$$

Then, this last equation, together with (7.9), provides

$$(7.11) \quad \sup_{z \in B(0, \frac{1}{\sqrt{2}})} \sum_{\gamma \neq e, \gamma \in \Gamma_g} (1 - |\gamma(z)|^2)^2 < \frac{1}{4}.$$

Remark 7.1. For the estimate (7.10) to hold, we need the ball, centered at the origin, to have Euclidean radius $\sqrt{\frac{99}{100}}$, so that the surface is at least $\ln \frac{10 + \sqrt{99}}{10 - \sqrt{99}}$ -thick.

The following two propositions follow easily from the equations above.

Proposition 7.2. *Let X_g be sufficiently thick. Then*

$$\sup_{z \in \mathcal{D}} |\mu_g(z)| < \frac{5}{16}.$$

Proof. We compute

$$\begin{aligned} |\mu_g(z)| &\leq \sum_{\gamma \in \Gamma_g} \frac{|\gamma'(z)|^2}{\rho(z)} \text{ by (7.1)} \\ &= \frac{1}{4} \sum_{\gamma \in \Gamma_g} (1 - |\gamma(z)|^2)^2 \text{ by (7.3)} \\ &\leq \frac{1}{4} \sup_{z \in \mathcal{D}} \sum_{\gamma \in \Gamma_g} (1 - |\gamma(z)|^2)^2 \\ &= \frac{1}{4} \sup_{z \in B(0, \frac{1}{\sqrt{2}})} \sum_{\gamma \in \Gamma_g} (1 - |\gamma(z)|^2)^2 \text{ by (7.5)} \\ &\leq \frac{1}{4} + \frac{1}{4} \sup_{z \in B(0, \frac{1}{\sqrt{2}})} \sum_{\gamma \neq e, \gamma \in \Gamma_g} (1 - |\gamma(z)|^2)^2 \\ &< \frac{5}{16} \text{ by (7.11)}. \end{aligned}$$

The conclusion follows by taking a supremum. \square

Proposition 7.3. *Let X_g be sufficiently thick as above. Then*

$$|\mu_g(0)| \geq \frac{3}{16}.$$

Proof. By the triangle inequality we have

$$\begin{aligned} |\mu_g(0)| &\geq \frac{1}{4} - \sum_{\gamma \neq e \in \Gamma_g} \frac{|\gamma'(0)|^2}{\rho(0)} \\ &= \frac{1}{4} - \frac{1}{4} \sum_{\gamma \neq e \in \Gamma_g} (1 - |\gamma(0)|^2)^2 \\ &> \frac{3}{16} \text{ by (7.11)}. \end{aligned}$$

\square

Remark 7.4. We have chosen our setting to be one of a single sufficiently thick surface. However, we could also consider a sequence $\{X_g\}$ of surfaces, say indexed by genus, whose injectivity radii $\text{inj}(X_g)$ is growing (without bound) with the genus g . In that case, the estimate (7.9) becomes

$$\sup_{z \in B(0, \frac{1}{\sqrt{2}})} \sum_{\gamma \neq e, \gamma \in \Gamma_g} (1 - |\gamma(z)|^2)^2 \leq \frac{25}{\pi} (\pi - \text{Area}(F_g)) = o_g(1)$$

so that the conclusions of the proofs of Propositions 1.1 and 7.3 become

$$|\mu_g(z)| \leq \frac{1}{4} + o_g(1)$$

and

$$|\mu_g(0)| \geq \frac{1}{4} - o_g(1)$$

respectively. We hope to take up the thread of the agreement of these asymptotic bounds in a subsequent work.

8. HOLOMORPHIC LINES WITH UNIFORM NEGATIVELY PINCHED WEIL-PETERSSON HOLOMORPHIC SECTIONAL CURVATURES.

We are now ready to prove Theorem 1.7, which we restate here in the present terminology for convenience.

Theorem 8.1. *Let X_g be a sufficiently thick surface as in section 7, and let $\mu_g \in T_{X_g} \text{Teich}(S_g)$ be defined as in equation (7.1). Then the holomorphic sectional curvature $K(\mu_g)$ satisfies*

$$K(\mu_g) \leq \frac{-81C_0}{6400 \cdot \pi^2} < 0$$

where $C_0 = \frac{2}{3C(1)^2 \cdot \text{Vol}_{\mathcal{D}}(B(0;1))}$.

Proof of Theorem 1.7. Recall $z = x + iy$. Our plan is to bound $\|\mu_g\|_{WP}^2$ from above, $\sup_{z \in X_g} |\mu_g(z)|$ from below and then apply Proposition 2.11 to bound the holomorphic sectional curvature. First,

(8.1)

$$\begin{aligned} \|\mu_g\|_{WP}^2 &= \int_{F_g} \mu_g \cdot \overline{\mu_g} \rho(z) dx dy \\ &\leq \sup_{z \in F_g} |\mu_g(z)| \cdot \int_{F_g} |\overline{\mu_g}| \rho(z) dx dy \\ &\leq \frac{5}{16} \int_{F_g} |\overline{\mu_g}| \rho(z) dx dy \text{ by Proposition 7.2, since } X_g \text{ is sufficiently thick} \\ &\leq \frac{5}{16} \int_{F_g} \sum_{\gamma \in \Gamma_g} |\gamma'(z)|^2 dx dy \text{ by (7.1)} \\ &\leq \frac{5}{16} \int_{\mathcal{D}} dx dy \\ &= \frac{5}{16} \pi. \end{aligned}$$

Second, by Proposition 7.3, again using that X_g is sufficiently thick,

$$\sup_{z \in X_g} |\mu_g(z)| = \sup_{z \in F_g} |\mu_g(z)| \geq |\mu_g(0)| \geq \frac{3}{16}.$$

Finally, we recall from Proposition 2.11 the bound

$$(8.2) \quad K(\mu_g) \leq -\frac{C_0 \cdot \sup_{z \in X} |\mu_g(z)|^4}{\|\mu_g\|_{WP}^4}.$$

Thus substituting (8.1) and (8.2) into the right hand side of (8.2), we find

$$K(\mu_g) \leq \frac{-81C_0}{6400 \cdot \pi^2},$$

as desired. \square

Our final task is to prove Corollary 1.8.

Proof of Corollary 1.8. We begin with the left hand side. For any $\sigma_g \in T_{X_g} \text{Teich}(S_g)$, Proposition 2.10 asserts that the normalized L^∞ norm of the Beltrami differential σ_g satisfies

$$\frac{\sup_{z \in X_g} |\sigma_g(z)|^2}{\int_{X_g} |\sigma_g(z)|^2 dA(z)} \leq C(\text{inj}(X_g))$$

where $C(\text{inj}(X_g))$ is the constant defined in (2.8). We then substitute this estimate into the bound in Proposition 2.11 to obtain the required lower bound. If, as we did at the end of section 7, we considered the setting of a sequence X_g of surfaces whose injectivity radii grow without bound, we could let $\text{inj}(X_g) \rightarrow \infty$ so that $C(\text{inj}(X_g)) \rightarrow \frac{3}{4\pi}$: we would then obtain the lower bound

$$-\frac{2}{\pi} \leq K(\sigma_g).$$

Since σ_g is arbitrary in $T_{X_g} \text{Teich}(S_g)$, the conclusion follows.

The upper bound is a direct consequence of the statement of the upper bound in Theorem 1.7, as that theorem only required sufficiently large thickness, which is immediately satisfied on our sequence X_g , once g is sufficiently large. \square

Remark 8.2. Fix two positive numbers a and b . Let $H(\mathcal{D})$ denote the holomorphic functions on the disk \mathcal{D} , and set

$$H_a^b(\mathcal{D}) = \{f(z) \in H(\mathcal{D}); \sup_{z \in \mathcal{D}} |f(z)| \leq b \text{ and } |f(0)| \geq a\}.$$

Let $f(z) \in H_a^b(\mathcal{D})$ and consider the image of the Theta-operator $\Theta(f)(z) \in H^0(X_g, K^2)$. Using the same argument in the proof of Theorem 1.7, we find

Theorem 8.3. *Let X_g be a sequence of hyperbolic surfaces with $\text{inj}(X_g) \rightarrow \infty$ as $g \rightarrow \infty$. Then there exists a uniform negative constant $B(a, b)$ such that for g large enough, the Weil-Petersson holomorphic sectional curvature satisfies*

$$\frac{-2}{\pi} \leq K(\mu_g(f)(z)) \leq B(a, b) < 0$$

where $\mu_g(f)(z) = \frac{\overline{\Theta(f)(z)}}{\rho(z)}$ and f is arbitrary in $H_a^b(\mathcal{D})$.

Remark 8.4. While our focus in this part of the paper was on the Beltrami differential μ_g defined in terms of the Θ -operator applied to the constant function $f \equiv 1$, we might equally well have applied the Θ -operator to the functions $f_n(z) = (1 - z^n)$ ($n \geq 1$) and obtained similar estimates since $f_n(z) \in H_1^2(\mathcal{D})$ for all $n \geq 1$. It is clear that the convex hull of $\{f_n(z)\}_{n \geq 1}$ is still contained in $H_1^2(\mathcal{D})$, It is then of interest to know the Weil-Petersson geometry of the image of this convex hull under the Θ -operator.

REFERENCES

- [1] Lars V. Ahlfors, *Some remarks on Teichmüller space of Riemann surfaces*, Ann. of Math. **74:2** (1961), 171-191.
- [2] Lars V. Ahlfors, *Eine Bemerkung über Fuchssche Gruppen*, Math. Z. **84** (1964), 244-245.
- [3] K. Burns, H. Masur and A. Wilkinson, *The Weil-Petersson geodesic flow is ergodic*, Ann. of Math. **175** (2012), 835-908.
- [4] P. Buser and P. Sarnak, *On the period matrix of a Riemann surface of large genus*, Invent. Math. **117** (1994), no. 1, 27-56. With an appendix by J. H. Conway and N. J. A. Sloane.
- [5] J. Brock, *The Weil-Petersson metric and volumes of 3-dimensional hyperbolic convex cores*, J. Amer. Math. Soc. **16:3** (2003), 495-535.
- [6] W. Cavendish and H. Parlier, *Growth of the Weil-Petersson diameter of moduli space*, Duke Math. J. **161:1** (2012), 139-171.
- [7] T. Chu, *The Weil-Petersson metric in the moduli space*, Chinese J. Math. **4:2** (1976), 29-51.
- [8] J. Hempel, *Residual finiteness of surface groups*, Proc. Amer. Math. Soc. **32** (1972), 323.
- [9] Z. Huang, *Asymptotic flatness of the Weil-Petersson metric on Teichmüller space*, Geom. Dedicata **110** (2005), 81-102.
- [10] Z. Huang, *The Weil-Petersson geometry on the thick part of the moduli space of Riemann surfaces*, Proc. Amer. Math. Soc. **135:10** (2007), 3309-3316 (electronic).
- [11] S. Lang, *Foundamentals of differential geometry*, Vol. **191** of Graduate Texts in Mathematics. Springer-Verlag, New York, 1999.
- [12] Y. Imayoshi and M. Taniguchi, *An introduction to Teichmüller spaces*, Springer-Verlag, Tokyo, 1992. 279pp.
- [13] J. Jost, *Harmonic maps and curvature computations in Teichmüller theory*, Ann. Acad. Sci. Fen., Series A. I. Math., **16** (1991), 13-46.
- [14] J. Jost, *Nonlinear methods in Riemannian and Kählerian geometry*, Second edition. DMV Seminar, 10. Birkhäuser Verlag, Basel, 1991. 154 pp.
- [15] K. Liu, X. Sun and S. T. Yau, *Good Geometry on the Curve Moduli*, Publ. Res. Inst. Math. Sci. **44:2** (2008), 699-724.
- [16] K. Liu, X. Sun and S. T. Yau, *Canonical metrics on the moduli space of Riemann surfaces*, I. J. Differential Geom. **68:3** (2004), 571-637.
- [17] K. Liu, X. Sun and S. T. Yau, *Canonical metrics on the moduli space of Riemann surfaces*, II. J. Differential Geom. **69** (2005), no. 1, 163-216.
- [18] K. Liu, X. Sun, X. Yang and S. T. Yau, *Curvatures of moduli space of curves and applications*, arXiv:1312.6932.
- [19] H. Masur, *The extension of the Weil-Petersson metric to the boundary of Teichmüller space*, Duke. Math. J. **43** (1976), 623-635.
- [20] D. Mumford, *A remark on Mahler's compactness theorem*, Proc. Amer. Math. Soc. **28** (1971), 289-294.

- [21] M. Mirzakhani, *Weil-Petersson volumes and intersection theory on the moduli space of curves*, J. Amer. Math. Soc. **20:1** (2007), 1-23.
- [22] M. Mirzakhani, *Simple geodesics and Weil-Petersson volumes of moduli spaces of bordered Riemann surfaces*. Invent. Math. **167** (2007), 179-222.
- [23] M. Mirzakhani, *Growth of the number of simple closed geodesics on hyperbolic surfaces*. Ann. of Math. **168** (2008), 97-125.
- [24] M. Mirzakhani, *Growth of Weil-Petersson volumes and random hyperbolic surfaces of large genus*. J. Differential Geom. **94** (2013), 267-300.
- [25] P. Petersen, *Riemannian geometry*, Vol. 171 of Graduate Texts in Mathematics. Springer, New York, second edition, 2006.
- [26] W. Roelcke, *Das Eigenwertproblem der automorphen Formen in der hyperbolischen Ebene*, I, II. (German) Math. Ann. **167** (1966), 292-337; *ibid.* **168** 1966 261-324.
- [27] G. Schumacher, *Harmonic maps of the moduli space of compact Riemann surfaces*, Math. Ann. **275:3** (1986) 455-466.
- [28] Y. T. Siu, *Curvature of the Weil-Petersson metric in the moduli space of compact Kähler-Einstein manifolds of negative first Chern class*. In *Contributions to several complex variables*, Aspects Math., E9, pages 261-298. Vieweg, Braunschweig, 1986.
- [29] L. P. Teo, *The Weil-Petersson geometry of the moduli space of Riemann surfaces*, Proc. Amer. Math. Soc. **137:2** (2009) 541-552.
- [30] A. J. Tromba, *On a natural algebraic affine connection on the space of almost complex structures and the curvature of Teichmüller space with respect to its Weil-Petersson metric*, Manuscripta Math. **56:4** (1986) 475-497.
- [31] M. Wolf, *The Teichmüller theory of harmonic maps*, J. Differential Geom. **29:2** (1989), 449-479.
- [32] M. Wolf, *The Weil-Petersson Hessian of length on Teichmüller space*, J. Differential Geom. **91:1** (2012), 129-169.
- [33] S. Wolpert, *Noncompleteness of the Weil-Petersson metric for Teichmüller space*, Pacific J. Math. **61:2** (1975), 573-577.
- [34] S. Wolpert, *Chern forms and the Riemann tensor for the moduli space of curves*, Invent. Math. **85** (1986:1), 119-145.
- [35] S. Wolpert, *Geodesic length functions and the Nielsen problem*, J. Differential Geom. **25** (1987), 275-296.
- [36] S. Wolpert, *Geometry of the Weil-Petersson completion of Teichmüller space*, Surveys in differential geometry, Vol. VIII (Boston, MA, 2002), 357-393, Surv. Differ. Geom., VIII, Int. Press, Somerville, MA, 2003.
- [37] S. Wolpert, *Behavior of geodesic-length functions on Teichmüller space*, J. Differential Geom. **79:2** (2008), 277-334.
- [38] S. Wolpert, *Families of Riemann surfaces and Weil-Petersson geometry*, CBMS Regional Conference Series in Mathematics, 113. Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 2010. viii+118 pp.
- [39] S. Wolpert, *Geodesic-length functions and the Weil-Petersson curvature tensor*, J. Differential Geom. **91:2** (2012), 321-359.
- [40] Y. Wu, *The Riemannian sectional curvature operator of the Weil-Petersson metric and its application*, J. Differential Geom. **96:3** (2014), 507-530.

DEPARTMENT OF MATHEMATICS, RICE UNIVERSITY, HOUSTON, TEXAS, 77005-1892,
E-mail address: mwolf@rice.edu

DEPARTMENT OF MATHEMATICS, RICE UNIVERSITY, HOUSTON, TEXAS, 77005-1892,
E-mail address: yw22@rice.edu