

# Math 211

Lecture #14

October 12, 2000

## Nullspace of a Matrix

The nullspace of a matrix  $A$  is the set

$$\text{null}(A) = \{\mathbf{x} \mid A\mathbf{x} = \mathbf{0}\}.$$

- The nullspace of  $A$  is the same as the solution set for the homogeneous system  $A\mathbf{x} = \mathbf{0}$ .
- What are the properties of nullspaces?
- Is there a convenient way to describe them?

return

## Subspaces of $\mathbf{R}^n$

**Definition:** A nonempty subset  $V$  of  $\mathbf{R}^n$  that has the properties

1. if  $\mathbf{x}$  and  $\mathbf{y}$  are vectors in  $V$ ,  $\mathbf{x} + \mathbf{y}$  is in  $V$ ,
2. if  $a$  is a scalar, and  $\mathbf{x}$  is in  $V$ , then  $a\mathbf{x}$  is in  $V$ ,

is called a subspace of  $\mathbf{R}^n$ .

- The nullspace of a matrix is a subspace.

return

## Examples of Subspaces

- The nullspace of a matrix is a subspace.
- A line through the origin is a subspace.  
 $V = \{t\mathbf{v} \mid t \in \mathbf{R}\}$ .
- A plane through the origin is a subspace.  
 $V = \{a\mathbf{v} + b\mathbf{w} \mid a, b \in \mathbf{R}\}$ .
- $\{0\}$  and  $\mathbf{R}^n$  are subspaces of  $\mathbf{R}^n$ .

return

## Linear Combinations

**Proposition:** Any linear combination of vectors in a subspace  $V$  is also in  $V$ .

- Subspaces of  $\mathbf{R}^n$  have the same kind of linear structure as  $\mathbf{R}^n$  itself.
- In particular the nullspaces of matrices have the same kind of linear structure as  $\mathbf{R}^n$ .

Outline

## Example of a Nullspace

$$A = \begin{pmatrix} 4 & 3 & -1 \\ -3 & -2 & 1 \\ 1 & 2 & 1 \end{pmatrix}$$

The nullspace of  $A$  is

$$\text{null}(A) = \{a\mathbf{v} \mid a \in \mathbf{R}\},$$

where  $\mathbf{v} = (1, -1, 1)^T$ .

return

## Example of a Nullspace

$$B = \begin{pmatrix} 4 & 3 & -1 & 6 \\ -3 & -2 & 1 & -4 \\ 1 & 2 & 1 & 4 \end{pmatrix}$$

- $\text{null}(B) = \{a\mathbf{v} + b\mathbf{w} \mid a, b \in \mathbf{R}\}$ , where  $\mathbf{v} = (1, -1, 1, 0)^T$  and  $\mathbf{w} = (0, -2, 0, 1)^T$ .
- $\text{null}(B)$  consists of all linear combinations of  $\mathbf{v}$  and  $\mathbf{w}$ .

return

## The Span of a Set of Vectors

In every example the subspace has been the set of all linear combinations of a few vectors.

**Definition:** The span of a set of vectors is the set of all linear combinations of those vectors.

The span of the vectors  $\mathbf{v}_1, \mathbf{v}_2, \dots$ , and  $\mathbf{v}_k$  is denoted by

$$\text{span}(\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k).$$

null(A)

null(B)

Examples

## The Span of a Set of Vectors

**Proposition:** If  $\mathbf{v}_1, \mathbf{v}_2, \dots$ , and  $\mathbf{v}_k$  are all vectors in  $\mathbf{R}^n$ , then  $V = \text{span}(\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k)$  is a subspace of  $\mathbf{R}^n$ .

Examples

Outline

## Examples

Let  $\mathbf{v}_1 = (1, 2)^T$ ,  $\mathbf{v}_2 = (1, 0)^T$ , and  $\mathbf{v}_3 = (2, 0)^T$ .

- $\text{span}(\mathbf{v}_1, \mathbf{v}_2) = \mathbf{R}^2$ . (Proof?)
- $\text{span}(\mathbf{v}_1, \mathbf{v}_3) = \mathbf{R}^2$ . (Proof?)
- $\text{span}(\mathbf{v}_2, \mathbf{v}_3) = \text{span}(\mathbf{v}_2)$ . (Proof?)
  - ◊  $\text{span}(\mathbf{v}_2, \mathbf{v}_3) = \{t\mathbf{v}_2 \mid t \in \mathbf{R}\}$ .
  - ◊  $\mathbf{v}_2$  and  $\mathbf{v}_3$  have the same direction.

return

## Linear Independence

We need a condition that will keep out superfluous vectors from a spanning list. We will work toward a definition.

- Two vectors are linearly dependent if one is a scalar multiple of the other.
  - ◊  $\mathbf{v}_2$  and  $\mathbf{v}_3$  are linearly dependent.
  - ◊  $\mathbf{v}_1$  and  $\mathbf{v}_2$  are linearly independent.

Examples

## Linear Independence

- Three vectors  $\mathbf{v}_1$ ,  $\mathbf{v}_2$ , and  $\mathbf{v}_3$  are linearly dependent if one is a linear combination of the other two.
  - ◊ Example:  $\mathbf{v}_1 = (1, 0, 0)^T$ ,  
 $\mathbf{v}_2 = (0, 1, 0)^T$ , and  $\mathbf{v}_3 = (1, 2, 0)^T$ 

$$\mathbf{v}_3 = \mathbf{v}_1 + 2\mathbf{v}_2.$$
  - ◊ Notice that  $\mathbf{v}_1 + 2\mathbf{v}_2 - \mathbf{v}_3 = \mathbf{0}$ .

return

## Linear Independence

- Three vectors are linearly dependent if there is a non-trivial linear combination of them which equals the zero vector.
  - ◊ Non-trivial means that at least one of the coefficients is not 0.
- A set of vectors is linearly dependent if there is a non-trivial linear combination of them which equals the zero vector.

LD3

## Linear Independence

**Definition:** The vectors  $\mathbf{v}_1, \mathbf{v}_2, \dots,$  and  $\mathbf{v}_k$  are linearly independent if the only linear combination of them which is equal to the zero vector is the one with all of the coefficients equal to 0.

- In symbols,

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \cdots + c_k\mathbf{v}_k = \mathbf{0}$$

$$\Rightarrow c_1 = c_2 = \cdots = c_k = 0.$$

return

## Basis of a Subspace

**Definition:** A set of vectors  $\mathbf{v}_1, \mathbf{v}_2, \dots,$  and  $\mathbf{v}_k$  form a basis of a subspace  $V$  if

1.  $V = \text{span}(\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k)$
2.  $\mathbf{v}_1, \mathbf{v}_2, \dots,$  and  $\mathbf{v}_k$  are linearly independent.

## Basis of a Subspace

- The vector  $\mathbf{v} = (1, -1, 1)^T$  is a basis for  $\text{null}(A)$ .
  - ◊  $\text{null}(A)$  is the subspace of  $\mathbf{R}^3$  with basis  $\mathbf{v}$ .
- The vectors  $\mathbf{v} = (1, -1, 1, 0)^T$  and  $\mathbf{w} = (0, -2, 0, 1)^T$  form a basis for  $\text{null}(B)$ .
  - ◊  $\text{null}(B)$  is the subspace of  $\mathbf{R}^4$  with basis  $\{\mathbf{v}, \mathbf{w}\}$ .

[null\(A\)](#)   [null\(B\)](#)   [Examples](#)   [Outline](#)   [return](#)

## Basis of a Subspace

**Proposition:** Let  $V$  be a subspace of  $\mathbf{R}^n$ .

1. If  $V \neq \{\mathbf{0}\}$ , then  $V$  has a basis.
2. Every basis of  $V$  has the same number of elements.

**Definition:** The dimension of a subspace  $V$  is the number of elements in a basis of  $V$ .

[Examples](#)

## Linear Independence?

How do we decide if a set of vectors is linearly independent?

$$\mathbf{v}_1 = \begin{pmatrix} 1 \\ -2 \\ 0 \\ 2 \end{pmatrix}, \quad \mathbf{v}_2 = \begin{pmatrix} -1 \\ -3 \\ 2 \\ 0 \end{pmatrix}, \quad \mathbf{v}_3 = \begin{pmatrix} 5 \\ 0 \\ -4 \\ 6 \end{pmatrix}$$

$$-3\mathbf{v}_1 + 2\mathbf{v}_2 + \mathbf{v}_3 = \mathbf{0}.$$

[Linear Independence](#)

$$\mathbf{v}_1 = \begin{pmatrix} 1 \\ -2 \\ 0 \\ 2 \end{pmatrix}, \mathbf{v}_2 = \begin{pmatrix} -1 \\ -3 \\ 2 \\ 0 \end{pmatrix}, \mathbf{v}_3 = \begin{pmatrix} 5 \\ 0 \\ -4 \\ 3 \end{pmatrix}$$

- These vectors are linearly independent.

Linear Independence

**Proposition:** Suppose that  $\mathbf{v}_1, \mathbf{v}_2, \dots,$  and  $\mathbf{v}_k$  are vectors in  $\mathbf{R}^n$ . Set  $V = [\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k]$ .

1. If  $\text{null}(V) = \{\mathbf{0}\}$ , then  $\mathbf{v}_1, \mathbf{v}_2, \dots,$  and  $\mathbf{v}_k$  are linearly independent.
2. If  $\mathbf{c} = (c_1, c_2, \dots, c_k)^T$  is a nonzero vector in  $\text{null}(V)$ , then

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_k\mathbf{v}_k = \mathbf{0},$$

so the vectors are linearly dependent.

Linear Independence

## Solutions Sets and Nullspaces

**Theorem:** The solution set of the inhomogeneous system  $A\mathbf{x} = \mathbf{b}$  has the form

$$\{\mathbf{x} = \mathbf{x}_p + \mathbf{v} \mid \mathbf{v} \in \text{null}(A)\},$$

where  $\mathbf{x}_p$  is any particular solution of the inhomogeneous system.

- Restatement of previous result.

## Determinants in 2D

- How do we decide if a matrix  $A$  is nonsingular?
- $A$  is nonsingular if and only if  $\text{rref}(A) = I$ .

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

is nonsingular if and only if  $ad - bc \neq 0$ .

- ◊ if and only if  $\det(A) = ad - bc \neq 0$

## Determinants in 3D

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}$$

$$\det(A) = a_{11}a_{22}a_{33} - a_{11}a_{23}a_{32} - a_{12}a_{21}a_{33} \\ + a_{12}a_{23}a_{31} - a_{13}a_{22}a_{31} + a_{13}a_{21}a_{32}.$$

- $A$  is nonsingular if and only if  $\det(A) \neq 0$ .

return

## Determinants

**Theorem:** The  $n \times n$  matrix  $A$  is nonsingular if and only if  $\det(A) \neq 0$ .

**Corollary:** If  $A$  is an  $n \times n$  matrix, then  $\text{null}(A)$  contains a nonzero vector if and only if  $\det(A) = 0$ .

- The corollary contains the most important fact about determinants for ODEs.

## Typical Term

- Typical term  $a_{13}a_{21}a_{32}$
- First subscripts are 1, 2, & 3. True for all terms.
- Second subscripts are 3, 1, & 2. Different order for each terms, but always a list of 1, 2, & 3.

3D

return

## Permutations

- A permutation is a list of the numbers 1, 2,  $\dots$ ,  $n$  in any order.
  - ◊ Example  $\sigma = (3, 1, 2)$ .
  - ◊ Typical term is  $a_{1\sigma_1}a_{2\sigma_2}a_{3\sigma_3}$
- All permutations of 3 numbers appear in the determinant of a  $3 \times 3$  matrix (sometimes with minus signs).

Typical term

3D

## Sign of a Permutation

Transform a permutation into  $(1, 2, \dots, n)$  using interchanges.

- $(3, 1, 2) \rightarrow (1, 3, 2) \rightarrow (1, 2, 3)$
- This can be done in many ways, but for a given permutation, the number of interchanges is always odd or always even.
- A permutation is odd if the number is odd, and even if the number is even.

## Sign of a Permutation

Define

$$(-1)^\sigma = \begin{cases} 1 & \text{if } \sigma \text{ is even,} \\ -1 & \text{if } \sigma \text{ is odd.} \end{cases}$$

## Definition of Determinant

The general  $n \times n$  matrix

$$A = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{pmatrix}$$

$$\det(A) = \sum_{\sigma} (-1)^\sigma a_{1\sigma_1} \cdot a_{2\sigma_2} \cdot \cdots \cdot a_{n\sigma_n}$$

return

## Definition of Determinant

- Each summand of the definition is the product of  $n$  entries, one from each row, and one from each column.
- The definition is the sum of  $n!$  terms. It does not provide an effective way to compute the determinant.
- One exception: The determinant of an upper triangular matrix is the product of the diagonal terms.

Definition

## Row Operations and Determinants

If  $B$  is obtained from  $A$  by

- adding a multiple of one row to another,  
 $\det(B) = \det(A)$ .
- interchanging two rows,  
 $\det(B) = -\det(A)$ .
- multiplying a row by  $c \neq 0$ ,  
 $\det(B) = c \det(A)$ .

return

