

Math 211

Lecture #1

August 29, 2000

Welcome to Math 211

Math 211 Section 4 – John C. Polking

Herman Brown 402
713-348-4841
polking@rice.edu

Office Hours: 1:30 – 2:30 TWTh

Ordinary Differential Equations with Linear Algebra

- Applications & modeling.
 - ◊ Mechanics, electric circuits, population genetics epidemiology, pollution, pharmacology, personal finance, etc.
- Analytic solutions.
- Numerical solutions.
- Qualitative analysis.
 - ◊ Properties of solutions without knowing what they are.

Math 211 Web Page

Official source of information about the course.

<http://www.owl.net.rice.edu/~math211/>.

What Is a Derivative?

- The rate of change of a function.
- The slope of the tangent line to the graph of a function.
- The best linear approximation to the function.
- The limit of difference quotients.
- Rules and tables that allow computation.

What Is an Integral?

- The area under the graph of a function.
- An anti-derivative.
- Rules and tables for computing.

Differential Equations

$$y' = f(t, y) \quad y' = 2ty$$

- t is the *independent variable*.
- y is the *unknown function*.
- This equation is of order 1.

Equations and Solutions

$$y' = f(t, y) \quad y' = 2ty$$

A solution is a function $y(t)$, defined for t in an interval, which is differentiable at each point and satisfies

$$y'(t) = f(t, y(t))$$

for every point t in the interval.

Example $y'(t) = 2ty(t)$.

Example: $y' = 2ty$

Claim: $y(t) = e^{t^2}$ is a solution.

Verify by substitution.

Left hand side: $y'(t) = 2te^{t^2}$

Right hand side: $2ty(t) = 2te^{t^2}$

Therefore $y' = 2ty$.

Types of Solutions

For the equation $y' = 2ty$

$y(t) = \frac{1}{2}e^{t^2}$ is a solution. It is a *particular solution*.

$y(t) = Ce^{t^2}$ is a solution for any constant C .

This is a *general solution*.

General solutions contain arbitrary constants.

Particular solutions do not.

Initial Value Problem

Consists of a differential equation and an initial condition.

E.g., $y' = -2ty$ and $y(0) = 4$.

General solution: $y(t) = Ce^{-t^2}$.

Initial condition:

$$y(0) = 4,$$

$$Ce^0 = 4,$$

$$C = 4$$

Solution to the IVP: $y(t) = 4e^{-t^2}$.

Normal Form of an Equation

$$y' = f(t, y)$$

Example:

$$(1 + t^2)y' + y^2 = t^3$$

Solve for y' to put into normal form:

$$y' = \frac{t^3 - y^2}{1 + t^2}$$

Interval of Existence

The largest interval over which a solution can exist.

Example: $y' = 1 + y^2$ with $y(0) = 1$

General solution: $y(t) = \tan(t + C)$

Initial Condition: $y(0) = 1 \Leftrightarrow C = \pi/4$.

Solution: $y(t) = \tan(t + \pi/4)$

$y(t)$ exists and is continuous for

$-\pi/2 < t + \pi/4 < \pi/2$

or for $-3\pi/4 < t < \pi/4$.

Geometric Interpretation of

$$y' = f(t, y)$$

If $y(t)$ is a solution, and $y(t_0) = y_0$, then

$$y'(t_0) = f(t_0, y(t_0)) = f(t_0, y_0).$$

The slope to the graph of $y(t)$ at the point (t_0, y_0) is given by $f(t_0, y_0)$.

Imagine a small line segment attached to each point of the (t, y) plane with the slope $f(t, y)$.

The Direction Field

