

Math 211

Lecture #25

November 28, 2000

Interacting Species

- Two species with populations x_1 & x_2 .
- Interaction between the species can be helpful or detrimental.
- Model

$$x'_1 = r_1 x_1$$

$$x'_2 = r_2 x_2$$

- r_1 & r_2 are the reproductive rates.

Return

Reproductive Rates

If $x_2 = 0$ the reproductive rate for x_1 is

$$r_1 = a_1 - b_1 x_1.$$

- $a_1 > 0 \Rightarrow$ natural growth.
- $a_1 < 0 \Rightarrow$ natural decline.
- $b_1 = 0$ Malthusian growth.
- $b_1 > 0$ logistic growth.

IS

Return

If $x_2 > 0$ the reproductive rate for x_1 is

$$r_1 = a_1 - b_1x_1 + c_1x_2.$$

- $c_1 > 0 \Rightarrow$ interaction is helpful to x_1 .
- $c_1 < 0 \Rightarrow$ interaction is detrimental to x_1 .

The reproduction rate for x_2 is

$$r_2 = a_2 - b_2x_2 + c_2x_1.$$

IS

RR1

Return

Interacting Species Model

$$x_1' = (a_1 - b_1x_1 + c_1x_2)x_1$$

$$x_2' = (a_2 - b_2x_2 + c_2x_1)x_2$$

Return

Predator Prey

Rabbits & foxes, fish & sharks, and cottony cushion scale insect & ladybird beetle.

- F = fish & S = sharks.

$$F' = (a - bS)F$$

$$S' = (-c + dF)S$$

or

$$F' = (a - eF - bS)F$$

$$S' = (-c + dF)S$$

ISM

RR1

RR2

Return

Example of Predator Prey

$$F' = (3 - 3S)F$$

$$S' = (-1 + 3F)S$$

or

$$F' = (3 - 3F - 3S)F$$

$$S' = (-1 + 3F)S$$

Return

Competing Species

Cattle and sheep.

- x_1 and x_2 competing for resources.

$$x_1' = (a_1 - b_1x_1 + c_1x_2)x_1$$

$$x_2' = (a_2 - b_2x_2 + c_2x_1)x_2$$

- $a_i > 0$, $b_i > 0$, & $c_i < 0$

ISM

RR1

RR2

Return

Example of Competing Species

$$x' = (5 - 2x - y)x$$

$$y' = (7 - 2x - 3y)y$$

CS

Return

Linearization

Principal idea of differential calculus:

- Approximate nonlinear mathematical objects by linear ones.
- Approximate $f(y) = f(y_0 + h)$ near y_0 by the linear function $L(h) = f(y_0) + f'(y_0)h$.

$$f(y_0 + h) = f(y_0) + f'(y_0)h + R(h)$$

$$\text{where } \lim_{h \rightarrow 0} \frac{R(h)}{h} = 0.$$

Return

Linearization of an ODE

$$y' = f(y)$$

- Assume $f(y_0) = 0$ and $f'(y_0) \neq 0$.
- Set $y = y_0 + u$. Get

$$u' = f'(y_0)u + R(u)$$

- Approximate by the linear equation

$$\tilde{u}' = f'(y_0)\tilde{u}$$

Der

Return

If $f'(y_0) \neq 0$ the equilibrium point of the linearization at 0 is the same as that of the nonlinear equation at y_0 .

- $f'(y_0) > 0 \Rightarrow y_0$ is unstable.
- $f'(y_0) < 0 \Rightarrow y_0$ is asymptotically stable.

Lin1

Return

Linearization of a Planar System

$$x' = f(x, y)$$

$$y' = g(x, y)$$

- (x_0, y_0) is an equilibrium point so

$$f(x_0, y_0) = g(x_0, y_0) = 0$$

Return

We have

$$\begin{aligned} f(x_0 + u, y_0 + v) \\ = \frac{\partial f}{\partial x}(x_0, y_0)u + \frac{\partial f}{\partial y}(x_0, y_0)v + R_f(u, v) \end{aligned}$$

$$\begin{aligned} g(x_0 + u, y_0 + v) \\ = \frac{\partial g}{\partial x}(x_0, y_0)u + \frac{\partial g}{\partial y}(x_0, y_0)v + R_g(u, v) \end{aligned}$$

$$\text{where } \frac{R_f(u, v)}{\sqrt{u^2 + v^2}} \rightarrow 0 \text{ and } \frac{R_g(u, v)}{\sqrt{u^2 + v^2}} \rightarrow 0$$

Return

- Set $x = x_0 + u$ and $y = y_0 + v$. The system becomes

$$u' = \frac{\partial f}{\partial x}(x_0, y_0)u + \frac{\partial f}{\partial y}(x_0, y_0)v + R_f(u, v)$$

$$v' = \frac{\partial g}{\partial x}(x_0, y_0)u + \frac{\partial g}{\partial y}(x_0, y_0)v + R_g(u, v)$$

Return

Linearization

$$\begin{aligned}\tilde{u}' &= \frac{\partial f}{\partial x}(x_0, y_0)\tilde{u} + \frac{\partial f}{\partial y}(x_0, y_0)\tilde{v} \\ \tilde{v}' &= \frac{\partial g}{\partial x}(x_0, y_0)\tilde{u} + \frac{\partial g}{\partial y}(x_0, y_0)\tilde{v}\end{aligned}$$

NLSYS

LSYSM

HDLIN

Return

Matrix Form of the Linearization

Set $\mathbf{u} = (\tilde{u}, \tilde{v})^T$ and introduce the Jacobian matrix

$$J = \begin{pmatrix} \frac{\partial f}{\partial x}(x_0, y_0) & \frac{\partial f}{\partial y}(x_0, y_0) \\ \frac{\partial g}{\partial x}(x_0, y_0) & \frac{\partial g}{\partial y}(x_0, y_0) \end{pmatrix}$$

- The linearization becomes

$$\mathbf{u}' = J\mathbf{u}.$$

LSYS

HDLIN

Return

Theorem: Consider the planar system

$$x' = f(x, y)$$

$$y' = g(x, y)$$

where f and g are continuously differentiable.

Suppose that (x_0, y_0) is an equilibrium point. If the linearization at (x_0, y_0) has a generic equilibrium point at the origin, then the equilibrium point at (x_0, y_0) is of the same type.

LSYS

LSYSM

Generic

Return

Generic Equilibrium Points

- Saddle, nodal source, nodal sink, spiral source, and spiral sink.
 - ◊ All occupy large open subsets of the trace-determinant plane.
- Nongeneric types
 - ◊ Center and eight others. Occupy pieces of the boundaries.

THM1

Return

Examples

- Predator prey
- Competing species
- Center

$$x' = y + \alpha x(x^2 + y^2)$$

$$y' = -x + \alpha y(x^2 + y^2)$$

THM1

Generic

Return