

Math 211

Lecture #25

Exponential Solutions

October 26, 2001

Homogeneous Systems

- These are systems of the form

$$\mathbf{x}' = A\mathbf{x},$$

where A is an $n \times n$ matrix.

- We are looking primarily at homogeneous systems with constant coefficients.

Structure of the Solution Space

Theorem: Suppose that $\mathbf{x}_1(t)$, $\mathbf{x}_2(t)$, \dots , and $\mathbf{x}_n(t)$ are linearly independent solutions to the $n \times n$ homogeneous system $\mathbf{x}' = A\mathbf{x}$ on the interval I . Then every solution is a linear combination of $\mathbf{x}_1(t)$, $\mathbf{x}_2(t)$, \dots , and $\mathbf{x}_n(t)$.

- That is, if $\mathbf{x}(t)$ is a solution, then there are constants C_1 , C_2 , \dots , and C_n such that

$$\mathbf{x}(t) = C_1\mathbf{x}_1(t) + C_2\mathbf{x}_2(t) + \dots + C_n\mathbf{x}_n(t).$$

Solution Strategy

- The obvious **strategy** for completely solving the system is to look for n linearly independent solutions.

Definition: A set of n linear independent solutions to the $n \times n$ homogeneous system $\mathbf{x}' = A\mathbf{x}$ is called a *fundamental set of solutions*.

- We will look for fundamental sets of solutions.

Procedure to Solve $\mathbf{x}' \equiv A\mathbf{x}$

- Find the **eigenvalues** of A
 - ◆ the roots of $p(\lambda) = \det(A - \lambda I) = 0$
- For each eigenvalue λ find the eigenspace
 - ◆ $= \text{null}(A - \lambda I)$
- If λ is an eigenvalue and \mathbf{v} is an associated eigenvector, $\mathbf{x}(t) = e^{\lambda t}\mathbf{v}$ is a solution.
- Hope that n of these are linearly independent.

Example

$$\mathbf{x}' = A\mathbf{x} \quad \text{where} \quad A = \begin{pmatrix} -4 & 2 \\ -3 & 1 \end{pmatrix}$$

- A has **eigenvalues** $\lambda_1 = -1$ and $\lambda_2 = -2$.
- Look for associated eigenvectors.
 - ♦ $\lambda_1 = -1$. Eigenvector: $\mathbf{v}_1 = (2, 3)^T$.
 - ▶ Solution: $\mathbf{x}_1(t) = e^{\lambda_1 t} \mathbf{v}_1 = e^{-t} (2, 3)^T$.
 - ♦ $\lambda_2 = -2$. Eigenvector: $\mathbf{v}_2 = (1, 1)^T$.
 - ▶ Solution: $\mathbf{x}_2(t) = e^{\lambda_2 t} \mathbf{v}_2 = e^{-2t} (1, 1)^T$.

- $\mathbf{x}_1(0) = \mathbf{v}_1 = (2, 3)^T$ and $\mathbf{x}_2(0) = \mathbf{v}_2 = (1, 1)^T$ are linearly independent.
- \mathbf{x}_1 and \mathbf{x}_2 form a fundamental set of solutions.
- The general solution is the set of all linear combinations:

$$\begin{aligned}\mathbf{x}(t) &= C_1\mathbf{x}_1(t) + C_2\mathbf{x}_2(t) \\ &= C_1e^{-t} \begin{pmatrix} 2 \\ 3 \end{pmatrix} + C_2e^{-2t} \begin{pmatrix} 1 \\ 1 \end{pmatrix}\end{aligned}$$

Solving $\mathbf{x}' = A\mathbf{x}$

Cases to be Considered

- Distinct real eigenvalues.
 - ◆ In this case the **method** works as described.
- Complex eigenvalues.
 - ◆ The method yields complex solutions, but we will want real solutions.
- Repeated eigenvalues.
 - ◆ The method does not always give enough solutions.
 - ▶ This is the hard case.

Planar System $\mathbf{x}' = A\mathbf{x}$

$$A = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \quad \text{and} \quad \mathbf{x}(t) = \begin{pmatrix} x_1(t) \\ x_2(t) \end{pmatrix}$$

In nonvector form

$$x_1' = a_{11}x_1 + a_{12}x_2$$

$$x_2' = a_{21}x_1 + a_{22}x_2$$

The Characteristic Polynomial

$$\begin{aligned} p(\lambda) &= \det(A - \lambda I) \\ &= \det \begin{pmatrix} a_{11} - \lambda & a_{12} \\ a_{21} & a_{22} - \lambda \end{pmatrix} \\ &= \lambda^2 - (a_{11} + a_{22})\lambda + (a_{11}a_{22} - a_{12}a_{21}) \\ &= \lambda^2 - T\lambda + D, \end{aligned}$$

where

- $D = a_{11}a_{22} - a_{12}a_{21} = \det(A)$
- $T = a_{11} + a_{22} = \text{tr}(A)$ is the *trace* of A .
 - ♦ The *trace* of a matrix is the sum of its diagonal elements.

The Eigenvalues of A

- The eigenvalues of A are the roots of the characteristic equation $p(\lambda) = \lambda^2 - T\lambda + D = 0$.

$$\lambda = \frac{T \pm \sqrt{T^2 - 4D}}{2}.$$

- Three cases:
 - ♦ 2 distinct real roots if $T^2 - 4D > 0$
 - ♦ 2 complex conjugate roots if $T^2 - 4D < 0$
 - ♦ Double real root if $T^2 - 4D = 0$

Eigenvectors are Linearly Independent

The problem of determining that solutions are linearly independent is eased by the following result.

Proposition: Suppose that $\lambda_1 \neq \lambda_2$ are eigenvalues of the $n \times n$ matrix A , and that $\mathbf{v}_1 \neq 0$ and $\mathbf{v}_2 \neq 0$ are eigenvectors associated with λ_1 and λ_2 , respectively. Then \mathbf{v}_1 and \mathbf{v}_2 are linearly independent.

Two Distinct Real Eigenvalues

$$\lambda_1 = \frac{T - \sqrt{T^2 - 4D}}{2}, \quad \lambda_2 = \frac{T + \sqrt{T^2 - 4D}}{2}$$

- $T^2 - 4D > 0$ so $\lambda_1 < \lambda_2$.
- There are associated nonzero eigenvectors \mathbf{v}_1 and \mathbf{v}_2 .
- Solutions $\mathbf{x}_1(t) = e^{\lambda_1 t} \mathbf{v}_1$ and $\mathbf{x}_2(t) = e^{\lambda_2 t} \mathbf{v}_2$.
- $\mathbf{x}_1(0) = \mathbf{v}_1$ and $\mathbf{x}_2(0) = \mathbf{v}_2$ are linearly independent ;
 $\mathbf{x}_1(t)$ and $\mathbf{x}_2(t)$ form a fundamental set of solutions.
- The general solution is $\mathbf{x}(t) = C_1 e^{\lambda_1 t} \mathbf{v}_1 + C_2 e^{\lambda_2 t} \mathbf{v}_2$.

Example

$$\mathbf{x}' = A\mathbf{x} \quad \text{where} \quad A = \begin{pmatrix} -6 & -8 \\ 4 & 6 \end{pmatrix}$$

- Characteristic polynomial: $p(\lambda) = \lambda^2 - 4$.
- Eigenvalues: $\lambda_1 = -2$ and $\lambda_2 = 2$.
 - ◆ $\lambda_1 = -2$. Eigenvector: $\mathbf{v}_1 = (-2, 1)^T$.
 - ▶ Solution: $\mathbf{x}_1(t) = e^{\lambda_1 t} \mathbf{v}_1 = e^{-2t} (-2, 1)^T$.
 - ◆ $\lambda_2 = 2$. Eigenvector: $\mathbf{v}_2 = (-1, 1)^T$.
 - ▶ Solution: $\mathbf{x}_2(t) = e^{\lambda_2 t} \mathbf{v}_2 = e^{2t} (-1, 1)^T$.

- \mathbf{x}_1 and \mathbf{x}_2 are a **fundamental set** of solutions.
- The general solution is

$$\begin{aligned}\mathbf{x}(t) &= C_1\mathbf{x}_1(t) + C_2\mathbf{x}_2(t) \\ &= C_1e^{-2t} \begin{pmatrix} -2 \\ 1 \end{pmatrix} + C_2e^{2t} \begin{pmatrix} -1 \\ 1 \end{pmatrix}.\end{aligned}$$

Initial Value Problem

Solve $\mathbf{x}' = A\mathbf{x}$ with the initial condition $\mathbf{x}(0) = (1, 4)^T$.

- We need

$$\mathbf{x}(0) = C_1\mathbf{x}_1(0) + C_2\mathbf{x}_2(0)$$

- ◆ $C_1 = -5$ and $C_2 = 9$.

- The solution is

$$\begin{aligned}\mathbf{x}(t) &= -5\mathbf{x}_1(t) + 9\mathbf{x}_2(t) \\ &= \begin{pmatrix} 10e^{-2t} - 9e^{2t} \\ -5e^{-2t} + 9e^{2t} \end{pmatrix}.\end{aligned}$$

Homogeneous Systems

$$\mathbf{x}' = A\mathbf{x}$$

Proposition: Suppose that $\mathbf{x}_1(t)$, $\mathbf{x}_2(t)$, \dots , and $\mathbf{x}_k(t)$ are solutions to the homogeneous system, and c_1, c_2, \dots , and c_k are scalars. Then

$$\mathbf{x}(t) = c_1\mathbf{x}_1(t) + c_2\mathbf{x}_2(t) + \dots + c_k\mathbf{x}_k(t)$$

is also a solution.

- Any linear combination of solutions to the homogeneous system is also a solution.

Linear Independence

Definition: A set of k solutions to the linear system $\mathbf{x}' = A\mathbf{x}$ is *linearly independent* if they are linearly independent at one value of t .

- Proposition \Rightarrow the solutions are linearly independent for all values of t .

Eigenvalues & Eigenvectors

Definition: λ is an *eigenvalue* of A if there is a nonzero vector \mathbf{v} such that $A\mathbf{v} = \lambda\mathbf{v}$. If λ is an eigenvalue of A , then any vector \mathbf{v} such that $A\mathbf{v} = \lambda\mathbf{v}$ is called an *eigenvector associated with λ* .

- λ an eigenvalue of A , \mathbf{v} an associated eigenvector
 $\Rightarrow \mathbf{x}(t) = e^{\lambda t}\mathbf{v}$ is a solution to $\mathbf{x}' = A\mathbf{x}$.
- The set of eigenvectors associated with the eigenvalue λ is the subspace $\text{null}(A - \lambda I)$, and is called the eigenspace of λ .