

# Math 211

Lecture #7

Mixing Problems

September 11, 2002

## Mixing Problem #1

A tank originally holds 500 gallons of pure water. At  $t = 0$  there starts a flow of sugar water into the tank with a concentration of  $\frac{1}{2}$  lbs/gal at a rate of 5 gal/min. There is also a pipe at the bottom of the tank removing 5 gal/min from the tank. Assume that the sugar is immediately and thoroughly mixed throughout the tank.

Find the amount of sugar in the tank after 10 minutes and after 2 hours.

# Model

- $S(t)$  = the amount of sugar in the tank in lbs.
- **Concentration** = pounds per unit volume.
  - ◆  $c(t) = \frac{S(t)}{V} \frac{\text{lbs}}{\text{gal}}$ .
- Modeling is easier in terms of the total amount,  $S(t)$ .
- Draw a picture.
- We must compute the rate of change of  $S$  in two ways.
  - ◆ The mathematical way: Rate of change =  $dS/dt$ .
  - ◆ The application way: This is where the real modeling takes place.

## The Rate of Change of $S(t)$

- Balance Law:

$$\text{Rate of change} = \text{Rate in} - \text{Rate out}$$

- Rate = volume rate  $\times$  concentration

- For the problem

$$\blacklozenge \text{ Rate in} = 5 \frac{\text{gal}}{\text{min}} \times \frac{1}{2} \frac{\text{lb}}{\text{gal}} = 2.5 \frac{\text{lb}}{\text{min}}$$

$$\blacklozenge \text{ Rate out} = 5 \frac{\text{gal}}{\text{min}} \times \frac{S}{500} \frac{\text{lb}}{\text{gal}} = \frac{S}{100} \frac{\text{lb}}{\text{min}}$$

## Solution

$$\begin{aligned}\frac{dS}{dt} &= \text{Rate in} - \text{Rate out} \\ &= 2.5 - \frac{S}{100}.\end{aligned}$$

- The equation is **linear**.
- General solution:  $S(t) = 250 + Ce^{-t/100}$ .
- **Particular solution**:  $S(t) = 250(1 - e^{-t/100})$ .

## Other possible initial conditions

- There is initially 20 lbs of sugar in the tank.
- The concentration of sugar in the tank at  $t = 0$  is 1 lb/gallon.

## Mixing Problem #2

A tank originally holds 500 gallons of sugar water with a concentration of  $\frac{1}{10}$  lb/gal. At  $t = 0$  there starts a flow of sugar water into the tank with a concentration of  $\frac{1}{2}$  lbs/gal at a rate of 5 gal/min. There is also a pipe at the bottom of the tank removing 10 gal/min from the tank. Assume that the sugar is immediately and thoroughly mixed throughout the tank.

Find the amount of sugar in the tank after 10 minutes and after 2 hours.

## Solution

- Rate in =  $5 \frac{\text{gal}}{\text{min}} \times \frac{1 \text{ lb}}{2 \text{ gal}} = 2.5 \frac{\text{lb}}{\text{min}}$
- Rate out =  $10 \frac{\text{gal}}{\text{min}} \times \frac{S \text{ lb}}{V \text{ gal}}$ 
  - ◆  $V(t) = 500 - 5t.$
  - ◆ Rate out =  $\frac{10S}{500 - 5t} \frac{\text{lb}}{\text{min}}$

$$\begin{aligned}\frac{dS}{dt} &= \text{Rate in} - \text{Rate out} \\ &= 2.5 - \frac{2S}{100 - t},\end{aligned}$$

- General solution:

$$S(t) = 2.5(100 - t) + C(100 - t)^2.$$

- Particular solution:

$$S(t) = 2.5(100 - t) - \frac{(100 - t)^2}{50}.$$

# Conjectures, Theorems, and Proof

- A *conjecture* is a statement that we think is true.
- A *theorem* is a statement for which we have a logical *proof*.
  - ◆ A theorem contains:
    - ▶ *hypotheses* (the assumptions made)
    - ▶ and *conclusions*
  - ◆ The conclusions are guaranteed to be true if the hypotheses are true.
  - ◆ The implication goes only one way.

## Example of a “Theorem”

If it rains the sidewalks get wet.

- Hypothesis — *If it rains*
- Conclusion — *the sidewalks get wet*

# Mathematics and Proof

- Theorems are proved by logical deduction.
- All of mathematics comes from a small number of very basic assumptions.
  - ◆ Called *axioms* or *postulates*.
- True of all parts of mathematics.
  - ◆ An algebraic derivation is an example of a proof.
- Definitions are not theorems.

# Solving the Linear Equation

$$x' = a(t)x + f(t)$$

Four step process:

1. Rewrite as  $x' - ax = f$ .
2. Multiply by the integrating factor

$$u(t) = e^{-\int a(t) dt}.$$

Equation becomes  $[ux]' = ux' - aux = uf$ .

3. Integrate:  $u(t)x(t) = \int u(t)f(t) dt + C$ .
4. Solve for  $x(t)$ .