

Math 211

Lecture #18

Properties of Solution Sets

October 7, 2002

Method of Solution for $Ax = b$

The method is called *elimination and backsolving*, or *Gaussian elimination*. There are four steps:

1. Use the augmented matrix $M = [A, b]$.
2. Use row operations to reduce the augmented matrix to row echelon form.
3. Write down the simplified system.
4. Backsolve.
 - ♦ Assign arbitrary values to the free variables.
 - ♦ Backsolve for the pivot variables.

Return

Elimination using MATLAB

- Row operations in MATLAB. $M = [A, b]$.
 - ♦ $R_i \rightarrow R_i + aR_j$
 - ▶ `>> M(i,:) = M(i,:) + a*M(j,:)`
 - ♦ $R_i \leftrightarrow R_j$
 - ▶ `>> M([i,j],:) = M([j,i],:)`
 - ♦ $R_i \rightarrow aR_i$
 - ▶ `>> M(i,:) = a*M(i,:)`
- The MATLAB command `rref`.

Return

Examples

$$A = \begin{pmatrix} -3 & 6 & 0 \\ -2 & 4 & 0 \\ -1 & 0 & 2 \end{pmatrix} \quad \mathbf{b}_1 = \begin{pmatrix} 2 \\ 3 \\ -5 \end{pmatrix} \quad \mathbf{b}_2 = \begin{pmatrix} -9 \\ -6 \\ 7 \end{pmatrix}$$

- The system $A\mathbf{y} = \mathbf{b}_1$ is inconsistent.
- The system $A\mathbf{y} = \mathbf{b}_2$ is consistent.

Return

Method

Consistent Systems

- A system is *consistent* if it has solutions.
- A system is consistent if and only if the simplified system is consistent.
- This is true if and only if the row echelon form of the augmented matrix does not have a pivot in its last column.

Return

Row echelon form

Examples

Homogeneous Systems

A homogeneous system has the form $A\mathbf{x} = \mathbf{0}$.

- The augmented matrix $M = [A, \mathbf{0}]$ has all zeros in the last column.
- During elimination the column of zeros is unchanged.
 - It is not really necessary to augment a homogeneous system.

Return

Homogeneous Systems (cont.)

- A homogeneous system is always consistent.
 - The zero vector $\mathbf{0}$ is always a solution.
- A homogeneous system $A\mathbf{x} = \mathbf{0}$ has a nonzero solution if and only if the row echelon form of A has a free column.
- A homogeneous system of n equations and m unknowns with $n < m$ always has a nonzero solution.
- A homogeneous system $A\mathbf{x} = \mathbf{0}$ where A is a square matrix has a nonzero solution if and only if there are vectors \mathbf{b} for which the system $A\mathbf{x} = \mathbf{b}$ has no solutions.

[Return](#)

Examples

$$A = \begin{pmatrix} 1 & 0 & 3 \\ 2 & -1 & 4 \\ 1 & 1 & 5 \end{pmatrix} \quad B = \begin{pmatrix} 1 & 0 & 3 \\ 2 & -1 & 4 \\ 1 & 1 & 4 \end{pmatrix}$$

[Return](#)

Singular and Nonsingular Matrices

The $n \times n$ matrix A is *nonsingular* if the equation $A\mathbf{x} = \mathbf{b}$ has a solution for any right hand side \mathbf{b} .

Proposition: The $n \times n$ matrix A is nonsingular if and only if the row echelon form of A has only nonzero entries along the diagonal.

- $\Leftrightarrow \text{rref}(A) = I$.

[Return](#)

[Examples](#)

Proposition: If the $n \times n$ matrix A is nonsingular then the equation $Ax = b$ has a *unique* solution for any right hand side b .

Proposition: The $n \times n$ matrix A is singular if and only if the homogeneous equation $Ax = 0$ has a non-zero solution.

- This is a result that we will use repeatedly.

Invertible Matrices

An $n \times n$ matrix A is *invertible* if there is an $n \times n$ matrix B such that $AB = BA = I$. The matrix B is called an *inverse* of A .

- If B_1 and B_2 are both inverses of A , then

$$B_1 = B_1(AB_2) = (B_1A)B_2 = B_2$$

- The inverse of A is denoted by A^{-1} .
- Invertible \Rightarrow nonsingular.
- Nonsingular \Rightarrow invertible.

Return

Computing the inverse A^{-1}

- Form the matrix $[A, I]$.
- Do elimination until the matrix has the form $[I, B]$.
- Then $A^{-1} = B$.

- Examples: $\begin{pmatrix} 1 & 2 \\ 2 & 3 \end{pmatrix}$, $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$, $B = \begin{pmatrix} 1 & 0 & 3 \\ 2 & -1 & 4 \\ 1 & 1 & 4 \end{pmatrix}$

- In MATLAB use `inv`.

Structure of the Solution Set

Theorem: Let \mathbf{x}_p be a particular solution to $A\mathbf{x}_p = \mathbf{b}$.

1. If $A\mathbf{x}_h = \mathbf{0}$ then $\mathbf{x} = \mathbf{x}_p + \mathbf{x}_h$ also satisfies $A\mathbf{x} = \mathbf{b}$.
 2. If $A\mathbf{x} = \mathbf{b}$, then there is a vector \mathbf{x}_h such that $A\mathbf{x}_h = \mathbf{0}$ and $\mathbf{x} = \mathbf{x}_p + \mathbf{x}_h$.
- Solution set for $A\mathbf{x} = \mathbf{b}$ is known if we know one particular solution \mathbf{x}_p and the solution set for the homogeneous system $A\mathbf{x}_h = \mathbf{0}$.

Return

Row operations

The permissible operations on the rows of the augmented matrix are called *row operations*.

- Add a multiple of one row to another.
- Interchange two rows.
- Multiply a row by a non-zero number.

Return

Row Echelon Form

A matrix is in *row echelon form* if every pivot lies strictly to the right of those in rows above.

$$\begin{pmatrix} P & * & * & * & * & * & * & * & * \\ 0 & P & * & * & * & * & * & * & * \\ 0 & 0 & 0 & P & * & * & * & * & * \\ 0 & 0 & 0 & 0 & P & * & * & * & * \\ 0 & 0 & 0 & 0 & 0 & 0 & P & * & * \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & P & * \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

- P is a pivot, $*$ is any number.

Return