

# Math 211

Lecture #28

Phase Plane Portraits

November 1, 2002

## Procedure to Solve $\mathbf{x}' = A\mathbf{x}$

- Find the eigenvalues of  $A$ , which are the roots of  $\det(A - \lambda I) = 0$ .
- For each eigenvalue  $\lambda$  find the eigenspace, which is equal to  $\text{null}(A - \lambda I)$ .
- If  $\lambda$  is an eigenvalue and  $\mathbf{v}$  is an associated nonzero eigenvector,  $\mathbf{x}(t) = e^{\lambda t}\mathbf{v}$  is a solution.
- Show that  $n$  of these are linearly independent, *if we can*.
  - ♦ This must be explored further if the system has dimension  $n \geq 3$ .

Return

## Planar System $\mathbf{x}' = A\mathbf{x}$

$$A = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \quad \text{and} \quad \mathbf{x}(t) = \begin{pmatrix} x_1(t) \\ x_2(t) \end{pmatrix}$$

- The characteristic polynomial is

$$p(\lambda) = \lambda^2 - T\lambda + D.$$

where

- ♦  $T = \text{tr } A = a_{11} + a_{22}$  and
- ♦  $D = \det A = a_{11}a_{22} - a_{12}a_{21}$ .

Return

Procedure

- The eigenvalues of  $A$  are the roots of  $p(\lambda) = \lambda^2 - T\lambda + D$ ,

$$\lambda = \frac{T \pm \sqrt{T^2 - 4D}}{2}.$$

- Three cases:
  - ♦ 2 distinct real roots if  $T^2 - 4D > 0$
  - ♦ 2 complex conjugate roots if  $T^2 - 4D < 0$
  - ♦ Double real root if  $T^2 - 4D = 0$

Return

## Real Distinct Eigenvalues

Suppose  $A$  is a real  $2 \times 2$  matrix with real eigenvalues  $\lambda_1 \neq \lambda_2$ , and associated nonzero eigenvectors  $\mathbf{v}_1$  and  $\mathbf{v}_2$ .

Then  $\mathbf{x}_1(t) = e^{\lambda_1 t} \mathbf{v}_1$  and  $\mathbf{x}_2(t) = e^{\lambda_2 t} \mathbf{v}_2$  form a fundamental set of solutions.

Return

Cases

## Complex Eigenvalues

Suppose  $A$  is a real  $2 \times 2$  matrix with complex conjugate eigenvalues  $\lambda$  and  $\bar{\lambda}$ , and associated nonzero eigenvectors  $\mathbf{w}$  and  $\bar{\mathbf{w}}$ .

Then

- $\mathbf{z}(t) = e^{\lambda t} \mathbf{w}$  and  $\bar{\mathbf{z}}(t) = e^{\bar{\lambda} t} \bar{\mathbf{w}}$  form a complex valued fundamental set of solutions, and
- $\mathbf{x}(t) = \operatorname{Re}(\mathbf{z}(t))$  and  $\mathbf{y}(t) = \operatorname{Im}(\mathbf{z}(t))$  form a real valued fundamental set of solutions.

Return

Cases

## The Degenerate Planar Case

- Find the (only) eigenvalue  $\lambda_1$ .
- Find an eigenvector  $\mathbf{v}_1 \neq \mathbf{0}$ .
- Find  $\mathbf{v}_2$  with  $(A - \lambda I)\mathbf{v}_2 = \mathbf{v}_1$ . To do so:
  - ♦ Start with any vector  $\mathbf{w}$  not a multiple of  $\mathbf{v}_1$
  - ♦ Then  $(A - \lambda I)\mathbf{w} = a\mathbf{v}_1$  with  $a \neq 0$ .
  - ♦ Set  $\mathbf{v}_2 = \frac{1}{a}\mathbf{w}$ .  $\mathbf{v}_2$  is not a multiple of  $\mathbf{v}_1$ .
- $\mathbf{x}_1(t) = e^{\lambda t}\mathbf{v}_1$  and  $\mathbf{x}_2(t) = e^{\lambda t}[\mathbf{v}_2 + t\mathbf{v}_1]$  form a fundamental set of solutions.

Return

Cases

## Qualitative Analysis for a Planar System

$$\mathbf{x}' = A\mathbf{x}$$

- Equilibrium points for the system
  - ♦ The set of equilibrium points equals  $\text{null}(A)$ .
  - ♦ If  $A$  is nonsingular, the only equilibrium point is  $\mathbf{0}$ .
- Can we list the types of all possible equilibrium points for planar linear systems?
  - ♦ The cases are distinguished by different solution curves in the phase plane.
  - ♦ We will do the six most important cases.

Return

Cases

## Distinct Real Eigenvalues

- $p(\lambda) = \lambda^2 - T\lambda + D$  with  $T^2 - 4D > 0$ .
- $$\lambda_1 = \frac{T - \sqrt{T^2 - 4D}}{2} < \lambda_2 = \frac{T + \sqrt{T^2 - 4D}}{2}$$

- Eigenvectors  $\mathbf{v}_1$  and  $\mathbf{v}_2$ . The general solution is

$$\mathbf{x}(t) = C_1 e^{\lambda_1 t} \mathbf{v}_1 + C_2 e^{\lambda_2 t} \mathbf{v}_2$$

- There are three subcases depending on the signs of the eigenvalues.

Return

Cases

Solution

## Exponential Solutions

$$\mathbf{x}(t) = Ce^{\lambda t}\mathbf{v}$$

- The solution curve is a straight half-line through  $C\mathbf{v}$ . Sometimes called *half-line* solutions.
- If  $\lambda > 0$  the solution starts at  $\mathbf{0}$  for  $t = -\infty$ , and tends to  $\infty$  as  $t \rightarrow \infty$ . This is an *unstable solution*
- If  $\lambda < 0$  the solution starts at  $\infty$  for  $t = -\infty$ , and tends to  $\mathbf{0}$  as  $t \rightarrow \infty$ . This is a *stable solution*

Return

Real case

## Saddle Point

- $\lambda_1 < 0 < \lambda_2$
- General solution  $\mathbf{x}(t) = C_1e^{\lambda_1 t}\mathbf{v}_1 + C_2e^{\lambda_2 t}\mathbf{v}_2$
- Two stable exponential solutions ( $C_2 = 0$ )
- Two unstable exponential solutions ( $C_1 = 0$ ).
- Suppose that  $C_1 \neq 0$  and  $C_2 \neq 0$ .
  - ♦ As  $t \rightarrow \infty$ ,  $\mathbf{x}(t) \rightarrow \infty$ , approaching the half-line through  $C_2\mathbf{v}_2$ .
  - ♦ As  $t \rightarrow -\infty$ ,  $\mathbf{x}(t) \rightarrow \infty$ , approaching the half-line through  $C_2\mathbf{v}_1$ .

Return

Real eigenvalues

## Nodal Sink

- $\lambda_1 < \lambda_2 < 0$
- General solution  $\mathbf{x}(t) = C_1e^{\lambda_1 t}\mathbf{v}_1 + C_2e^{\lambda_2 t}\mathbf{v}_2$
- Four stable exponential solutions.
- All solutions  $\rightarrow \mathbf{0}$  as  $t \rightarrow \infty$ . (Stable)
  - ♦ Tangent to  $C_2\mathbf{v}_2$  if  $C_2 \neq 0$ .
- All solutions  $\rightarrow \infty$  as  $t \rightarrow -\infty$ .
  - ♦  $\parallel$  to the half line through  $C_1\mathbf{v}_1$  if  $C_1 \neq 0$ .

Return

Real eigenvalues

## Nodal Source

- $0 < \lambda_1 < \lambda_2$
- General solution  $\mathbf{x}(t) = C_1 e^{\lambda_1 t} \mathbf{v}_1 + C_2 e^{\lambda_2 t} \mathbf{v}_2$
- Four unstable exponential solutions.
- All solutions  $\rightarrow \mathbf{0}$  as  $t \rightarrow -\infty$ .
  - ♦ Tangent to  $C_1 \mathbf{v}_1$  if  $C_1 \neq 0$ .
- All solutions  $\rightarrow \infty$  as  $t \rightarrow \infty$ . (Unstable)
  - ♦  $\parallel$  to the half line through  $C_2 \mathbf{v}_2$  if  $C_2 \neq 0$ .

Return

Real eigenvalues

Nodal Sink

## Complex Eigenvalues

- $p(\lambda) = \lambda^2 - T\lambda + D$  with  $T^2 - 4D < 0$   
 $\lambda = \alpha + i\beta$  and  $\bar{\lambda} = \alpha - i\beta$ .
- Eigenvector  $\mathbf{w} = \mathbf{v}_1 + i\mathbf{v}_2$  associated to  $\lambda$ .
- Complex solutions

$$\mathbf{z}(t) = e^{\lambda t} \mathbf{w} = e^{t(\alpha+i\beta)} [\mathbf{v}_1 + i\mathbf{v}_2]$$

$$\bar{\mathbf{z}}(t) = e^{\bar{\lambda} t} \bar{\mathbf{w}} = e^{t(\alpha-i\beta)} [\mathbf{v}_1 - i\mathbf{v}_2]$$

Return

Cases

Solution

- Real solutions
 
$$\mathbf{x}_1(t) = \operatorname{Re}(\mathbf{z}(t)) = e^{\alpha t} [\cos \beta t \cdot \mathbf{v}_1 - \sin \beta t \cdot \mathbf{v}_2]$$

$$\mathbf{x}_2(t) = \operatorname{Im}(\mathbf{z}(t)) = e^{\alpha t} [\sin \beta t \cdot \mathbf{v}_1 + \cos \beta t \cdot \mathbf{v}_2]$$
- General solution
 
$$\mathbf{x}(t) = C_1 e^{\alpha t} [\cos \beta t \cdot \mathbf{v}_1 - \sin \beta t \cdot \mathbf{v}_2]$$

$$+ C_2 e^{\alpha t} [\sin \beta t \cdot \mathbf{v}_1 + \cos \beta t \cdot \mathbf{v}_2]$$
- There are three cases depending on the sign of  $\alpha = \operatorname{Re}(\lambda)$ .

Return

Cases

Solution

## Center

- $\alpha = \operatorname{Re}(\lambda) = 0$
- The general real solution is

$$\mathbf{x}(t) = C_1[\cos \beta t \cdot \mathbf{v}_1 - \sin \beta t \cdot \mathbf{v}_2] \\ + C_2[\sin \beta t \cdot \mathbf{v}_1 + \cos \beta t \cdot \mathbf{v}_2]$$

- Every solution is periodic with period  $T = 2\pi/\beta$ .
- All solution curves are ellipses.

Return

## Spiral Sink

- $\alpha = \operatorname{Re}(\lambda) < 0$
- The general real solution is

$$\mathbf{x}(t) = C_1 e^{\alpha t}[\cos \beta t \cdot \mathbf{v}_1 - \sin \beta t \cdot \mathbf{v}_2] \\ + C_2 e^{\alpha t}[\sin \beta t \cdot \mathbf{v}_1 + \cos \beta t \cdot \mathbf{v}_2]$$

- All solutions spiral into  $\mathbf{0}$  as  $t \rightarrow \infty$ .

Return

## Spiral Source

- $\alpha = \operatorname{Re}(\lambda) > 0$
- The general real solution is

$$\mathbf{x}(t) = C_1 e^{\alpha t}[\cos \beta t \cdot \mathbf{v}_1 - \sin \beta t \cdot \mathbf{v}_2] \\ + C_2 e^{\alpha t}[\sin \beta t \cdot \mathbf{v}_1 + \cos \beta t \cdot \mathbf{v}_2]$$

- All solutions spiral into  $\mathbf{0}$  as  $t \rightarrow -\infty$ .

Return

## Planar Systems

$$A = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}$$

- The characteristic polynomial is  $p(\lambda) = \lambda^2 - T\lambda + D$ .  
where
  - ♦  $T = \text{tr } A = a_{11} + a_{22}$  and
  - ♦  $D = \det A = a_{11}a_{22} - a_{12}a_{21}$ .
- The eigenvalues are

$$\lambda_1, \lambda_2 = \frac{T \pm \sqrt{T^2 - 4D}}{2}.$$

Return

Planar systems

- $\lambda_1$  &  $\lambda_2$  are the roots of  $p(\lambda) = \lambda^2 - T\lambda + D$ , so

$$\begin{aligned} p(\lambda) &= (\lambda - \lambda_1)(\lambda - \lambda_2) \\ &= \lambda^2 - (\lambda_1 + \lambda_2)\lambda + \lambda_1\lambda_2 \end{aligned}$$

- Hence,  $T = \lambda_1 + \lambda_2$  and  $D = \lambda_1\lambda_2$ .
- Duality between  $(\lambda_1, \lambda_2)$  and  $(T, D)$ .
- We will represent a system by the location of  $(T, D)$  in the  $TD$ -plane — the *trace-determinant plane*.

Return

## Trace-Determinant Plane

- $T^2 - 4D > 0$ 
  - ♦  $\Rightarrow$  distinct real eigenvalues  $\lambda_1$  &  $\lambda_2$
  - ♦  $D = \lambda_1\lambda_2 < 0 \Rightarrow$  Saddle point.
  - ♦  $D = \lambda_1\lambda_2 > 0 \Rightarrow$  Eigenvalues have the same sign.
    - $T = \lambda_1 + \lambda_2 > 0 \Rightarrow$  Nodal source.
    - $T = \lambda_1 + \lambda_2 < 0 \Rightarrow$  Nodal sink.

Return

Duality

- $T^2 - 4D < 0 \Rightarrow$  complex eigenvalues

$$\lambda = \alpha + i\beta \quad \text{and} \quad \bar{\lambda} = \alpha - i\beta.$$

- ♦  $T = \lambda + \bar{\lambda} = 2\alpha > 0 \Rightarrow$  Spiral source.
- ♦  $T = \lambda + \bar{\lambda} = 2\alpha < 0 \Rightarrow$  Spiral sink.
- ♦  $T = \lambda + \bar{\lambda} = 2\alpha = 0 \Rightarrow$  Center.

[Return](#)

[Duality](#)

[TD plane](#)

[Cases](#)

## Types of Equilibrium Points

- *Generic* types
  - ♦ Saddle, nodal source, nodal sink, spiral source, and spiral sink.
  - ♦ All occupy large open subsets of the trace-determinant plane.
- *Nongeneric* types
  - ♦ Center and many others. Occupy pieces of the boundaries between the generic types.

[Return](#)