

Math 211

Lecture #7

Mixing Problems

September 10, 2003

Mixing Problem #1

A tank originally holds 500 gallons of pure water. At $t = 0$ there starts a flow of sugar water into the tank with a concentration of $\frac{1}{2}$ lbs/gal at a rate of 5 gal/min. There is also a pipe at the bottom of the tank removing 5 gal/min from the tank. Assume that the sugar is immediately and thoroughly mixed throughout the tank.

Find the amount of sugar in the tank after 10 minutes and after 2 hours.

Model

- $S(t)$ = the amount of sugar in the tank in lbs.
- **Concentration** = pounds per unit volume.
 - ◆ $c(t) = \frac{S(t)}{V} \frac{\text{lbs}}{\text{gal}}$.
- Modeling is easier in terms of the total amount, $S(t)$.
- Draw a picture.
- We must compute the rate of change of S in two ways.
 - ◆ The mathematical way: Rate of change = dS/dt .
 - ◆ The application way: This is where the real modeling takes place.

The Rate of Change of $S(t)$

- Balance Law:

$$\text{Rate of change} = \text{Rate in} - \text{Rate out}$$

- Rate = volume rate \times concentration
- For the problem

$$\blacklozenge \text{ Rate in} = 5 \frac{\text{gal}}{\text{min}} \times \frac{1}{2} \frac{\text{lb}}{\text{gal}} = 2.5 \frac{\text{lb}}{\text{min}}$$

$$\blacklozenge \text{ Rate out} = 5 \frac{\text{gal}}{\text{min}} \times \frac{S}{500} \frac{\text{lb}}{\text{gal}} = \frac{S}{100} \frac{\text{lb}}{\text{min}}$$

- The model equation is

$$\frac{dS}{dt} = 2.5 - \frac{S}{100}.$$

Solution

$$\frac{dS}{dt} = 2.5 - \frac{S}{100}$$

- The equation is **linear**.
- General solution: $S(t) = 250 + Ce^{-t/100}$.
- **Particular solution**: $S(t) = 250(1 - e^{-t/100})$.
- Other possible initial conditions
 - ♦ There is initially 20 lbs of sugar in the tank.
 - ♦ The concentration of sugar in the tank at $t = 0$ is 1 lb/gallon.

Mixing Problem #2

A tank originally holds 500 gallons of sugar water with a concentration of $\frac{1}{10}$ lb/gal. At $t = 0$ there starts a flow of sugar water into the tank with a concentration of $\frac{1}{2}$ lbs/gal at a rate of 5 gal/min. There is also a pipe at the bottom of the tank removing 10 gal/min from the tank. Assume that the sugar is immediately and thoroughly mixed throughout the tank.

Find the amount of sugar in the tank after 10 minutes and after 2 hours.

Solution

- Rate in = $5 \frac{\text{gal}}{\text{min}} \times \frac{1}{2} \frac{\text{lb}}{\text{gal}} = 2.5 \frac{\text{lb}}{\text{min}}$

- Rate out = $10 \frac{\text{gal}}{\text{min}} \times \frac{S}{V} \frac{\text{lb}}{\text{gal}}$

- ♦ $V(t) = 500 - 5t$, so Rate out = $\frac{10S}{500 - 5t} \frac{\text{lb}}{\text{min}}$

- The model equation is

$$\frac{dS}{dt} = \text{Rate in} - \text{Rate out} = 2.5 - \frac{2S}{100 - t}.$$

- **General solution:** $S(t) = 2.5(100 - t) + C(100 - t)^2.$

- **Particular solution:** $S(t) = 2.5(100 - t) - \frac{(100 - t)^2}{50}.$

Conjectures, Theorems, and Proof

- A *conjecture* is a statement that we think is true.
- A *theorem* is a statement for which we have a logical *proof*.
 - ◆ A theorem contains:
 - ▶ *hypotheses* (the assumptions made)
 - ▶ and *conclusions*
 - ◆ The conclusions are guaranteed to be true if the hypotheses are true.
 - ◆ The implication goes only one way.

Example of a “Theorem”

If it rains the sidewalks get wet.

- Hypothesis — *If it rains*
- Conclusion — *the sidewalks get wet*

Mathematics and Proof

- Theorems are proved by logical deduction.
- All of mathematics comes from a small number of very basic assumptions.
 - ◆ Called *axioms* or *postulates*.
- True of all parts of mathematics.
 - ◆ An algebraic derivation is an example of a proof.
- Definitions are not theorems.

Solving Linear Equations

To solve $x' = a(t)x + f(t)$:

1. Solve the homogeneous equation $x'_0 = ax_0$.
2. Find v such that $x = vx_0$ is a solution by substituting into the equation.
3. Write down the general solution, $x(t) = v(t)x_0(t)$.