

# Math 211

Lecture #8

Existence & Uniqueness

September 12, 2003

## Qualitative Analysis

- Do solutions always exist?
  - ♦ Do solutions to an initial value problem always exist?
- How many solutions are there?
  - ♦ How many solutions are there to an initial value problem?
- If we solve an IVP with an initial condition that is slightly wrong will the computed solution be close to the real one?
- Can we predict the behavior of solutions without having a formula?

[Return](#)

## Existence Theorem

**Theorem:** Suppose the function  $f(t, y)$  is defined and continuous in the rectangle  $R$  in the  $ty$ -plane. Then given any point  $(t_0, y_0) \in R$ , the initial value problem

$$y' = f(t, y) \quad \text{with} \quad y(t_0) = y_0$$

has a solution  $y(t)$  defined in an interval containing  $t_0$ . Furthermore the solution will be defined at least until the solution curve  $t \rightarrow (t, y(t))$  leaves the rectangle  $R$ .

[Return](#)

### Explanation of the Existence Theorem

- Hypotheses:
  - ♦ The equation is in normal form  $y' = f(t, y)$ .
  - ♦ The right hand side,  $f(t, y)$ , is continuous in the rectangle  $R$ .
  - ♦ The initial point  $(t_0, y_0)$  is in the rectangle  $R$ .
- Conclusions:
  - ♦ There is a solution starting at the initial point.
  - ♦ The solution is defined at least until the solution curve  $t \rightarrow (t, y(t))$  leaves the rectangle  $R$ .

[Return](#)

### Existence of a Solution

- The existence theorem does not guarantee an explicitly defined solution.
- In the proof, the solution is constructed as the limit of a sequence of explicitly defined functions.
- Frequently no explicit formula is possible.
- An ordinary differential equation is a function generator.

### Interval of Existence

- Example:  $y' = 1 + y^2$  with  $y(0) = 0$ .
- RHS  $f(t, y) = 1 + y^2$  is defined and continuous on the whole  $ty$ -plane. The rectangle  $R$  can be any rectangle in the plane.
- Solution  $y(t) = \tan t$  "blows up" at  $t = \pm\pi/2$ .
- Thus the size of the rectangle on which  $f(t, y)$  is continuous does not say much about the interval of existence.

[Existence theorem](#)

[Return](#)

## Uniqueness of Solutions

- How many solutions does an initial value problem have?
- The uniqueness of solutions to an initial value problem is the mathematical equivalent of being able to predict results in science and engineering.
- The uniqueness of solutions to a differential equation model is equivalent to a system being causal.

Questions

## Example of Non-uniqueness

- Initial value problem

$$y' = y^{1/3} \quad \text{with} \quad y(0) = 0.$$

- The constant function  $y_1(t) = 0$  is a solution.
- Solve by separation of variables to find that

$$y_2(t) = \begin{cases} \left(\frac{2t}{3}\right)^{3/2}, & \text{if } t > 0 \\ 0 & \text{if } t \leq 0. \end{cases}$$

is also a solution.

Return

Existence theorem

## Uniqueness Theorem

**Theorem:** Suppose the function  $f(t, y)$  and its partial derivative  $\partial f / \partial y$  are continuous in the rectangle  $R$  in the  $ty$ -plane. Suppose that  $(t_0, x_0) \in R$ . Suppose that

$$x' = f(t, x) \quad \text{and} \quad y' = f(t, y),$$

and that

$$x(t_0) = y(t_0) = x_0.$$

Then as long as  $(t, x(t))$  and  $(t, y(t))$  stay in  $R$  we have

$$x(t) = y(t).$$

Return

Example

Existence theorem

### Uniqueness Theorem

- Hypotheses:
  - ♦ The equation is in normal form  $y' = f(t, y)$ .
  - ♦ The right hand side,  $f(t, y)$ , and its derivative  $\partial f/\partial y$  are continuous in the rectangle  $R$ .
  - ♦ The initial point  $(t_0, y_0)$  is in the rectangle  $R$ .
- Conclusions:
  - ♦ There is one and only one solution starting at the initial point.
  - ♦ The solution is defined at least until the solution curve  $t \rightarrow (t, y(t))$  leaves the rectangle  $R$ .

Return

### Geometric Interpretation

- Solution curves cannot cross.
- They cannot even touch at one point.
- $y' = (y - 1)(\cos t - y)$  and  $y(0) = 2$ . Show that  $y(t) > 1$  for all  $t$ .
- $y' = y - (1 - t)^2$  and  $y(0) = 0$ . Show that  $y(t) < 1 + t^2$  for all  $t$ .

Uniqueness theorem

Hypotheses and Conclusions

### E & U for Linear Equations

**Theorem:** Suppose that  $a(t)$  and  $g(t)$  are continuous on an interval  $I = (a, b)$ . Then given  $t_0 \in I$  and any  $y_0$ , the initial value problem

$$y' = a(t)y + g(t) \quad \text{with} \quad y(t_0) = y_0$$

has a unique solution  $y(t)$  which exists for all  $t \in I$ .

- Notice that the RHS is

$$f(t, y) = a(t)y + g(t), \quad \text{and} \quad \frac{\partial f}{\partial y} = a(t).$$

These are continuous for  $t \in I$  and all  $y$ .

Existence theorem

Uniqueness theorem

## DFIELD6

Get a geometric look at existence and uniqueness.

**Theorem:** Suppose  $f(t, y)$ ,  $\partial f/\partial y$  are continuous in the rectangle  $R$ . Let

$$M = \max_{(t,y) \in R} \left| \frac{\partial f}{\partial y}(t, y) \right|.$$

Suppose that  $(t_0, x_0)$  and  $(t_0, y_0)$  both lie in  $R$ , and

$$x' = f(t, x), \quad x(t_0) = x_0 \quad \&$$

$$y' = f(t, y), \quad y(t_0) = y_0.$$

Then as long as  $(t, x(t))$  and  $(t, y(t))$  stay in  $R$  we have

$$|x(t) - y(t)| \leq |x_0 - y_0| e^{M|t-t_0|}.$$

[Return](#)

[Uniqueness theorem](#)

## Continuity in Initial Conditions

- Inequality:  $|x(t) - y(t)| \leq |x_0 - y_0| e^{M|t-t_0|}$ .
- The good news:
  - ♦ By making sure that  $x_0$  and  $y_0$  are very close we can make the solutions  $x(t)$  and  $y(t)$  close for  $t$  in an interval containing  $t_0$ .
  - ♦ Solutions are *continuous in the initial conditions*.

[Return](#)

### Sensitivity with Respect to Initial Conditions

- Inequality:  $|x(t) - y(t)| \leq |x_0 - y_0|e^{M|t-t_0|}$ .
- The bad news:
  - ♦ As  $|t - t_0|$  increases the RHS grows exponentially.
  - ♦ Over long intervals in  $t$  the solutions can get very far apart. Solutions are *sensitive to initial conditions*.

Return

Good news