

# Math 211

Lecture #42

Immunity to Infectious Diseases

December 5, 2003

## Model of the Immune System

- How does the immune system develop immunity to virus caused diseases?
  - ♦ Such as flu, the cold, mumps, . . .
- *Infectious Diseases of Humans* - Roy M. Anderson & Robert M. May, Oxford University Press 1992
- The model includes the interactions between virus cells and lymphocytes generated by the immune system.
  - ♦  $V(t)$  = number of virus cells
  - ♦ Two types of lymphocytes,  $E_1(t)$  &  $E_2(t)$ .

Return

## The Lymphocytes

- Both types of lymphocytes:
  - ♦ are recruited from bone marrow at a constant rate
  - ♦ die at a rate proportional to their numbers
  - ♦ proliferate due to contact with each other
- The model with no virus present is:


$$E_1' = \Lambda_1 - \mu_1 E_1 + a_1 \frac{E_1 E_2}{1 + b_1 E_1 E_2}$$

$$E_2' = \Lambda_2 - \mu_2 E_2 + a_2 \frac{E_1 E_2}{1 + b_2 E_1 E_2}$$

- ♦ In `pp1ane6` use parameters  $\Lambda_1 = \Lambda_2 = 1$ ,  $\mu_1 = \mu_2 = 1.25$ ,  $a_1 = a_2 = 0.252$ , and  $b_1 = b_2 = 0.008$ .

Return

## The Dynamics of the Lymphocytes



Return

## Interactions with the Virus

- Virus cells have an intrinsic growth rate  $r$ .
- Lymphocytes of type  $E_1$ :
  - ♦ kill virus because of contacts with them
  - ♦ proliferate because of contacts with virus
- Lymphocytes of type  $E_2$ :
  - ♦ do not directly interact with the virus
  - ♦ regulate cells of type  $E_1$

Return

## The Model With the Virus Present

$$E_1' = \Lambda_1 - \mu_1 E_1 + a_1 \frac{E_1 E_2}{1 + b_1 E_1 E_2} + K V E_1$$

$$E_2' = \Lambda_2 - \mu_2 E_2 + a_2 \frac{E_1 E_2}{1 + b_2 E_1 E_2}$$

$$V' = r V - k V E_1$$

Return

No virus

Interactions

### Analysis of the System

- There are three realistic equilibrium points

$$\begin{pmatrix} E_1 \\ E_2 \\ V \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 5 \\ 5 \\ 0 \end{pmatrix}, \quad \& \quad \begin{pmatrix} 20 \\ 20 \\ 0 \end{pmatrix}$$

- The first two are unstable. The third is asymptotically stable.
- What is the global behavior?
  - ♦ The best we can do is to compute with ode45. (Use  $K = 0.5$ ,  $k = 0.01$  and  $r = 0.1$ .)

Return

No virus